

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

NEW PLANTS

Sent out for the first time by the Company in 1883.



The prices are quoted in shillings (S) and pence (D).



Begonia amabilis.

This new species resembles slightly the varieties of *B. Teuscheri* which we sent out last year and which met with such a brilliant success. The leaves are of a bronze green speckled with silver, the underside being of a claret red contrasts favourably with the bright markings of the upperside. Native of Sumatra.

Price : 10s.

Begonia Meysselliana.

This species, with pale green leaves maculed with silver, is, as well as the preceding, very suitable for the open-air during summer. It is specially suitable for bordering purposes. Native of Sumatra.

Price : 5 s.

Curcuma Leopoldi.

Very pretty plant resembling a *Musa*, with lanceolate leaves, pale green, striped with creamy white. It grows in clumps and forms a very distinct and attractive stove plant.

Price : 20 s.

Calamus Lindenii




The *Calamus Lindenii* is one of the most striking Palms introduced up to this day. The leaves are thick, of a brilliant green on the upperside and on the under of a beautiful white. Its habit is like that of a Calamus or *Acanthophœnix*, but no species of these two classes approaches it in beauty.

We unfortunately only possess a very limited stock which we offer our honourable clients, in young plants at :

125s. each.


Dieffenbachia magnifica



E. VERHORCKEN sc.

The *Dieffenbachia magnifica* was introduced last year from Venezuela by one of our collectors. It is a remarkable species, with large obtuse leaves, of a clear-green marked with white like the *D. seguine picta* but also with a few yellow and green blotches intermingled forming a veritable mosaic where all the different shades of white and green are to be found. In fact it is one of the most beautiful Dieffenbachias ever introduced.

Price) Nice Young Plant : 30s.
) Half Specimen : 63s.



Nephrodium Rodriguezianum.

Impatiens Sultani

A most lovely novelty, native of central Africa. It is rather dwarf with bright carmine flowers which are produced in its earliest state on the tops of the stems in thick clusters and which form a most attractive head of flowers.

It will be a most useful plant during the summer for bedding out purposes as its propagation is as rapid and easy as that of the coleus, Iresine etc.

Price : 2s.

5s. the 3 plants.

Nephrodium Rodigasianum.

Illustration Horticole, Feby 1882.

Introduced from the Samoan isles, this remarkable fern was first exposed at the international Flower-Show at Antwerp in 1881 where it created a great surprise on account of the graceful hanging of its foliage.

The engraving will give a slight idea of its beauty.

Price : 25s.

Pellionia Daveauana var. *argentea*.

This variety introduced from Cochinchina differs from the original in having the upper side of the leaves, the stems and the petioles slightly velvety and the leaves are sometimes pure green, sometimes marked with silver.

Price : 4s.

Pellionia pulchra. N. E. Brown.

Illustration Horticole. March 1883.

Among the plants for hanging which exist in our collections the *Pellionias*, especially this variety, must occupy a prominent position.

Introduced, recently from Cochinchina this new species resembles in habit the *P. Daveauana*, but it differs from this latter in the form and colour of the leaves.

They are elongated with the upper surface green whereas the midrib and veins are blackish which produces a charming effect.

Besides, the leaves are not shining as are those of the *P. Daveauana*, but they have an opaque and slightly velvety appearance.

Price : 4s.

Panax fruticosum var. *Deleauana*. N. E. Brown.

The lovely series of Panax has been enriched by one other graceful species with the leaves divided like an Adiantum; it is native of the Salomon Isles from whence we received it last year. The leaves are of a lovely tender green deeply indented and resembling the leaves of the *Adiantum Luddemanianum*.

Price : 20 s.

Schismatoglottis Lansbergeana

L'Illustration Horticole. Nov. 1882.



A splendid novelty which will certainly find a great success among ornamental foliage plant collections.

The chief difference in this new variety is the colour of the upper side of the leaves which are of a dark green, whereas the stems and the under side of the leaves are of a claret color. The contrast is most delightful, and we feel sure this new aroid will receive a hearty welcome from amateurs of this fine group of plants.

Price : 20s.
Show Plant : 100s.

NEW SONERILAS

L'ILLUSTRATION Horticole. September 1882.

The 6 varieties offered here were selected from several hundred hybrids as being the best and most distinct.

Sonerila Comtesse de Flandre.

This variety resembles the *S. Mamei* but it is a great improvement on this latter; the petioles are much redder and the margin of the leaf is also lightly tinted with the same shade.

Sonerila Madame Alfred Mame.

Foliage with a groundwork of silver speckled with green points the veins being of a deeper hue. One of the best varieties.

Sonerila Madame Charles Heine.

Leaves of a pale green with deep-colored veining and numerous hieroglyphic markings.

Sonerila Madame Legrelle.

This lovely novelty resembles the *S. M^{me} Ed. Otlet*, only the green veining is much deeper.

Sonerila Madame Secretan.

The stems and petioles are reddish, as well as the under side of the young leaves. Markings of a pearly hue are scattered irregularly on the upper side.

Sonerila Princesse Mathilde.


It is an improvement of the *S. Hendersoni argentea*, which we sent out in 1876. The plant is much shorter in habit. The edge of the leaves is dark brown pointed with pea green on a white surface.

The six novelties will, we feel sure, be welcomed by all who are amateurs of this lovely little group of gems.

Each single variety : 10s.
The six varieties : 50s.

Spathiphyllum hybridum N. E. Brown.

L'Illustration Horticole. May 1882.



This lovely novelty is a hybrid obtained by us between the *S. Patini* and the *S. cannaefolium* (*Anthurium Dechartria*).

In foliage it resembles the *S. cannaefolium*, but its spathe, of a pure white on either side, differs from both its parents. It will be a most desirable acquisition for the confection of bouquets during the winter at which period it flowers abundantly.

Price : 25s.

Half specimen : 75s.

Show specimen : 150s.

Vriesea heliconoides. Ed. Morren.

Sent out under the name of *Vriesea bellula*.

It is one of the prettiest Bromeliads known. The leaves of a metallic purple surround the flower spike, which consists of bracts half carmin and white enclosing little white flowers. The habit is at the same time most graceful.

Price : 25s.


Vriesea Rodigasiana

L'Illustration Horticole. November 1882.



Native of southern Brazil this Bromeliad is one of the most brilliant we have ever seen. The eminent authority on these lovely plants, M. Ed. Morren, in describing it speaks of it in the highest terms. After such praise it is quite useless to add any other notes. Only a very few are available.

Price : 50s.



Aglaonema pictum.

NEW PLANTS

sent out for the first time by the Company on the 1st May 1882

Aglaonema Pictum, Kunth.

Illustration Horticole. March 1882.

The *Aglaonema pictum*, is a brilliant Aroid for the stove, with a fine rounded stem. The leaves vary in form, sometimes elliptical, sometimes oblong or oblong-lanceolate, are rounded at the base and almost suddenly acuminate at the top, from 4 to 7 inches long, and 2 to 2 3/4 in width; they are of a lovely deep velvety green, beautifully mottled with large spots and irregular markings of a whitish hue. The mid-rib is somewhat prominent and slightly channelled on the upper face, prominent and rounded beneath; with 4 or 6 principal lateral veins on each side, which are sunken on the upper surface and protruding beneath. When it is in flower, its white inflorescence makes it still more attractive. The spathe is remarkable for its curious pocket-like form and its knotty spadix visible at the top of the orifice.

Price : 15 s.

Alocasia Putzeysi, Brown.

Illustration Horticole. January 1882.

The general appearance of this plant resembles that of the *Alocasia longiloba*, but from which it is easily distinguished by the brilliant colouring of the foliage, which gives it a highly decorative value; in short, this magnificent variety is not surpassed in beauty by any of its predecessors. The *Alocasia Putzeysi* was discovered in Sumatra and is dedicated to the memory of Jules Putzeys, formerly general secretary to the Ministry of Justice at Brussels, a great promoter of botany.

Aralia Reginæ, Lind.

Illustration Horticole. February 1879.

This very ornamental species belongs to the section which has already furnished us with so many graceful and fashionable miniatures, such as *Aralia Veitchi, elegantissima, filifera*.

But in the variety we offer to-day, the even stem and the rose and olive petioles are marked with deeper freckles, and the palmate divisions of the stalk are fine, smooth, and of a uniform green. This plant was much admired at all the large exhibitions during the last three years.

A few specimens only are disposable.

Price : 50s.

Araucaria Mülleri, Ad. Brong. et Gris

Illustration Horticole. April 1882.



It is a very remarkable species with long pinnules, gracefully pendant, of a rich reddish-brown colour, having the same tint as the leaves of *Phormium-atro-purpureum*.

Its culture and hardiness are the same as that of the other greenhouse Araucarias.

This will be a valuable acquisition for the gardens on the Mediterranean.

Young rooted cutting : 25s.

Begonia diadema, Lind.

Illustration horticole, March 1882.

It is without dispute a species of incomparable beauty and moreover of a distinct type from all those which have hitherto been classed in the Begonia family. From the short and fleshy stem, branch out numerous leaves seated on rounded petioles, erect, of a delicate green, and marked with white elongated blotches. The leaves are deeply digitate-lobed; the lobes are irregular, acuminate-ovate, with the margin irregularly dentated; and are perfectly smooth and shiny. The ground is a clear green, whereon numerous white stains are scattered, without regularity, in bands or longitudinal series contained between the secondary veins. Little dentations mark the centre of each spot.

Price : 3s and 5s.

Begonia Teuscheri, Lind.

Illustration horticole, 1879.

- 1. **Begonia Teuscheri**
- 2. — — — **var. punctata**
- 3. — — — **var. maculata**

The *B. Teuscheri* is quite a new type, and is especially meritorious as it is excellent for cultivating in the open air during the warm season. The texture of the leaf is thick, fleshy, brittle, with a purple margin, unequally, dentated like a saw.

The variety : 4s.

The three " : 10s.

Caraguata Cardinalis, Ed. And.

Illustration horticole, 1880.

The beautiful scarlet bracts of this species, are of an intense brilliancy and preserve their colours for a long time.

The heads, of those plants which have flowered in our establishment, have developed with great vigour.

They shewed golden bracts forming the center of this brilliant inflorescence, which gave the appearance of a compact head of flowers, with white twisted petals, slightly projecting, buried in a thick mucilage as is meet with in other Tillandsias and also in several Orchids belonging to the Evelynia genus.

Young plant from seed : 50s.

Strong plant for flowering issuing from division : 100s.

Only a few examples are ready.

Croton magnificum, Lind.

Illustration horticole, 1882.

The variety we have named *magnificum*, well deserves the title, as it is one of the best and most distinct. The leaves are long, average size, of a clear green shaded with bands of bright yellow and purple.

The old leaves are entirely variegated with this colour, whereas the young ones are almost entirely yellow. This will be one of the best varieties and one which gives the greatest contrast.

Price : 15s.

Hibiscus rosa-sinensis Lucien Linden, Brown

Illustration horticole, January 1882.



This splendid ornamental shrub, is without doubt one of the best varieties of the kind. The flowers are large and very double, of a pale yellow colour beautifully streaked with spots and lines of brilliant red.

Introduced into our establishment since 1881, we have remarked with pleasure the rapid advance of the *Hibiscus Rosa-sinensis* in the favour of the public.

This world-wide praise is well deserved when we consider the graceful contrast it produces, when grouped with other fine varieties such as the lovely *H. Rosa-sinensis puniceus*.

Price : 10s.

Heliconia Triumphans, Lind.

Illustration Horticole. April 1882



Very ornamental plant with bright green leaves, lined with greenishblack. It is a robust species having the port of a *Maranta zebra* with shining foliage and which succeeds well in apartments. It is native of Sumatra. At the expositions at Brussels and Ghent, during last summer, it was much admired and the connoisseurs predicted a successful future for it.

Price : 10s.

Strong plant : 50s.

Piper Metallicum, Lind.

Very decorative plant with rounded leaves, thick and of a beautiful sparkling metallic green. This very distinct species is native of Borneo.

It is a meritorious plant and which ought to be in every stove. We recommend it with confidence.

Price : 10s.

 The collection of 13 new plants for 1882 are available at 140s. and 360s. according to force.

NEW PLANTS

*Sent out for the first time by the Establishment on
the 1st April 1881.*

GREAT REDUCTION IN PRICE.

The two collections of 12 new plants, exhibited by the Establishment at the spring Exhibition of 1881 of the SOCIÉTÉ ROYALE D'AGRICULTURE ET DE BOTANIQUE at Ghent, obtained respectively first and second certificates.

Schismatoglottis (*Aglaonema*) Lavallee, L.

Illustration horticole, 1881.

The stems are somewhat slender, cylindrical, and slightly knotted between the petioles; the leaf-blades of a clear green clasp the stem deeply. The leaf blade is elongate-lanceolate, obliquely heart-shaped at the base. The texture of the leaf is flexible; the mid-rib is slightly prominent on the upper surface, and of a beautiful bright green, on which are scattered numerous si'very white spots, producing a most pleasing effect.

Price : 10s.

Fine plant : 25s. and 50s.

Amorphophallus Lacourii

For figure and description, see Illustration horticole, n° 6, 1878.

A plant, the habit resembling that of its congeners, but distinguished by the form of the segments of the leaves and their white spotting which is similar to several of the Dieffenbachias. It is of fine aspect, and certainly the most beautiful of the genus.

Fine bulb : Price 10s. and 25s.

Aspidium Germinyi, Lind.

Illustration horticole, n° 12, 1880.

Nothing could be fresher or more graceful than this charming new Fern that was so much admired at the National Exhibition of Brussels last year, where it figured in the collection of six new plants, with which we obtained the First Prize. The fronds grow from 20 in. to 40 in. in length, elegantly recurved and terminated, as well as each pinnule, by a kind of leafy tuft.

Native of the South Sea Isles, it requires a cold house and does well in rooms.

Price : 3s. and 10s.

Exhibition plant : 25s. and 50s.

Clivia (Imantophyllum) Lindeni, Hort.

Illustration horticole, 1879.


Among last year's Novelties the *C. Lindeni* is without doubt the most brilliant variety. All those who had admired the beautiful specimens which flowered last year and in the present spring in the houses of the Count Adrien de Germiny à Gouville, or in those of the Count de Kerchove de Denterghem, and of M. Maurice de Ghellinck de Walle, of Ghent, are strongly convinced that it will be difficult, if not impossible, to surpass its large and brilliantly-coloured clusters of flowers. We have no hesitation in predicting for *C. Lindeni* a brilliant success, and in considering it as the perfection of the genus.

Price, young plants : 20s. 50s.

Some strong plants are available.

Colocasia Neo-Guineensis, Lind.

For figure and description, see Illustration Horticole. n° 5, 1880.



A new Aroid, introduced from New Guinea. It is remarkable for its tufted habit, the shortness of the leaf stalks, and its short-stalked inflorescence, and

the beautiful green tone of its smooth and shiny leaves, spotted with creamy white. It will form an interesting addition to collections of Aroidaceous plants in our hothouses.

Price : 5s. and 10s.

Strong plant : 25s.

Dracæna Lindeni, Hort. Lind.

Figure and description in Illustration Horticole. n° 6, 1880.

One of the most brilliant acquisitions recently made among fine-foliaged plants and a worthy rival of *D. Goldieana*.

We are fortunate to be able to offer to the public a form which, while it is in no way inferior to its forerunners in beauty, is distinct from them by its habit and pleasing aspect. The stem of *D. Lindeni* is erect and simple; the leaves are arranged regularly, deeply stem clasping at the base; the petiole is channelled, and winged, afterwards widening into a lanceolate acuminate blade, margined by white and rose and copiously traversed by regular bands of white and pale yellow, shaded and intermixed with zones of green, with others white in the middle. It is this particular arrangement, the blending of these two coloured bands, that gives the special merit to this superior form of Dracæna. It is a plant which will doubtless gain a high reputation, but we fear it will unfortunately be distributed but slowly, especially for the impatience of amateurs, as its propagation is very difficult.

The *Dracæna Lindeni* is a plant for a cool house where it colours far more brilliantly than in a hotter house.

Price : 50s.

Strong plants : 100s. and 250s.


Licuala (Pritchardia) Grandis, Wendland

Illustration Horticole. n° 2, 1881.

This is a palm of the highest order, the introduction of which has been eagerly desired by lovers of this beautiful family. We are happy to announce that we have had the good fortune to introduce a certain number of young plants, which arrived in excellent condition. The numerous visitors to the International Horticultural Exhibition of Brussels in 1876, Paris and Versailles in 1878, will no doubt remember the admiration this brilliant novelty caused when shown there under the name of *Pritchardia grandis*. It is this same species which we now offer.

Price : 125s. each.

Pothos aurea.



Pothos aurea, Linden.

For figure and description, see Illustration Horticole. n° 5, 1880.

A valuable introduction for the embellishment of rockeries in hothouses and winter gardens. This first class novelty is especially remarkable for the abundance of its much branched creeping and climbing stems, and by the golden-yellow colour which variegates its foliage in a very irregular, yet extremely elegant manner. It is a native of the Solomon Isles, and is amenable to the treatment accorded to Dracænas and Crotons, which for several years past have been introduced from these regions to adorn our hothouses.

Price : 2s. and 5s.

Strong plant : 10s. and 25s.

Selaginella Vogeli, Foliis variegatis

Among the Selaginellas with variegated foliage, this one is the most in demand as much on account of its vigorous habit as for its brilliant aspect.

Price : 3s. and 5s.



NEW "BEDDING-OUT" PLANT

Gynura Aurantiaca, Brown.

The *Illustration horticole* for 1881, last part, speaks thus of this beautiful plant.

"The *Gynura aurantiaca* is a hardy plant belonging to the Compositae, and is of such an ornamental character as to allow of one's saying that it is not surpassed by any other plant of the same class. The stem and leaves are clothed throughout their entire length with a thick covering of hairs, soft to the touch, and of a beautiful deep violet colour, which gives an appearance of the richest velvet to the plant. This is more especially the case with the young leaves, and when combined with the brilliant orange of the flowers, the aspect of the plant is truly superb."

Since its introduction in 1882 this plant has had a brilliant success, every where it is largely employed for bedding-out and under the effects of a hot summer forms a most brilliant and striking effect.

Price : 2s.

PALMS.

The Establishment possesses the richest and completest collection of Palms in the trade.

Those varieties which are of greenhouse cultivation, or suitable for rooms, halls, etc. as well as for the open air in the south of Europe, are marked with an †.

CHOICE OF THE BEST VARIETIES.

Acanthophœnix Alexandræ		Astrocaryum tenuifolium 25s.
(<i>Ptychosperma</i>)	1-5s.	Attalea compta	5s.
— Herbsti	10-25s.	Bentinckia Condapanna	
— rubra (<i>Calamus dealbatus et Verschaffeltii</i>) . .	10-25s.	(<i>New</i>)	10s.
Acanthorhiza aculeata (<i>Chamaerops stauracantha</i>)		†Brahea calcarea 25s.
(<i>Illustr.</i>)	10-25s.	— dulcis (<i>Illustr.</i>)	15s.
Areca alba (<i>Dictyosperma</i>) 2-10s.	Calamus asperrimus (<i>Illustr.</i>)	3-10s.
— Aliaeae (<i>new</i>)	25s.	— calilepis	25s.
— aurea (<i>Dictyosperma</i>) . .	3-15s.	— ciliaris	25s.
† — Baueri (<i>Seaforthia robusta</i>)	2-15s.	† Lewisianus (<i>Illustr.</i>)	5-15s.
— Betel	5-15s.	— Lindeni (<i>New</i>)	125s.
— lutescens (<i>Hyophorbe indica</i>)	1-2-5s.	— nitidus	15-30s.
— madagascariensis	2-10s.	— sumatranaus	10-25s.
† — monostachya.	2-10s.	— zeylanicus	15-30s.
— nobilis	5-50s.	Calyptrogyne elata 10-25s.
— purpurea (<i>Illustr.</i>) . .	3-5-10s.	Caryota elegans 10s.
— rubra (<i>Dictyosperma</i>) . .	2-5s.	— excelsa	10s.
† — sapida (<i>Rhopalostylis</i>) . .	2-10s.	— majestica.	15s.
— Verschaffeltii (<i>Illustr.</i>)		— Rumphi	15s.
(<i>Hyophorbe Verschaffeltii</i>)	3-10s.	— sobolifera (<i>vera</i>)	5-15s.
Arenga saccharifera (Sugar palm) 2-25s.	— speciosa	15-25s.
Astrocaryum argenteum 15s.	— Teysmanniana	10-25s.
— Ayri	10-25s.	— urens (<i>Illustr.</i>)	3-10s.
— Malybo	25s.	† Ceroxylon nivale (<i>Dipterium caudescens</i>) 10-50s.
— mexicanum	15-25s.	Chamædorea amazonica 10s.
— Murumuru (<i>Illustr.</i>) . .	25s.	— elegans	5-100s.
		— elegantissima	5-10s.
		— Ernesti-Augusti (<i>simplificifrons</i>)	10s.
		— graminifolia.	10-25s.
		— Lindeniana	5s.

Chamædorea lunata	10-50s.	† Cocos Romanzoffiana	2-20s.
— <i>pygmæa</i>	100s.	— Weddelliana (Illustr.)	3-25s.
— <i>Wendlandi</i>	10-25s.	— <i>de Yurumaguas</i>	30s.
Chamærops arborea	5s.	† Corypha (Livistona) australis	1-5s.
† — <i>elegans</i>	10-25s.	† — <i>Gebanga</i>	5s.
† — <i>excelsa vera</i>	1-10s.	Cyrtostachys Renda (new)	50s.
† — <i>Fortunei (sinensis)</i>	1-10s.	Dæmonorops cinnamomeus	10-25s.
† — <i>humilis</i>	1-10s.	— <i>fissus</i>	10-25s.
† — — <i>robusta</i>	5-10s.	— hystrix	25-50s.
† — <i>tomentosa</i>	5-20s.	— <i>marginatus</i>	10-25s.
† — hystrix (true)	25s.	— <i>ornatus</i>	20s.
† — <i>Martiana</i>	15s.	— <i>palembanicus</i>	10-25s.
† — <i>Palmetto</i>	2-10s.	— <i>periacanthus</i>	10s.
— <i>stauracantha (Acantho-</i> <i>rhiza)</i>	10-25s.	— <i>platiacanthus</i>	10s.
Cocos australis (<i>Diplothemium</i> <i>campestre</i>)	2-10s.	— <i>spectabilis</i>	25s.
† — <i>Blumenaui</i>	5-15s.	Deckenia corneto	25s.
† — <i>Bonneti</i>	5-15s.	Desmoncus granatensis	25s.
† — <i>campestris</i>	5-25s.	— <i>sp nova</i>	20s.
† — <i>chilensis (Jubaea spec-</i> <i>tabilis)</i>	2-10s.	— <i>mexicanus</i>	25s.
† — <i>flexuosa</i>	3-25s.	† Diplothemium caudescens	
— <i>Maria Regia</i>	30-50s.	(<i>Ceroxylon nivaleum</i>)	10-50s.
— <i>Maximiliana</i>	10s.	Drymophlaeus Rumphi	20s.
† — <i>Mikaniana</i>	25-50s.	Dypsis elata	15s.
— <i>nucifera</i>	30s.	Elæis guineensis (Illustr.)	10-25s.
— <i>oleracea</i>	5-15s.	Euterpe antioquensis (mon-	
† — <i>plumosa</i>	10-25s.	<i>tana</i>)	5-10s.
		— <i>edulis</i>	1-5s.
		— <i>elegans</i>	5-10s.
		— <i>montana</i>	25s.
		— <i>oleracea</i>	3-15s.
		Geonomia antioquensis	10-25s.
		— <i>elegans</i>	15-25s.
		— <i>fenestralis (Malortica)</i>	25s.
		— <i>gracilis</i>	3-5-10s.
		— <i>princeps</i>	5-10s.
		— <i>pumila</i>	15s.
		— <i>Seemannii</i>	15-25s.
		— <i>speciosa</i>	15s.
		Hyophorbe amaricaulis . (Voir <i>areca speciosa</i>)	3-10s.
		— <i>elata</i>	10s.
		— <i>indica (Areca lutescens)</i>	1-5s.
		— <i>Verschaffeltii</i>	3-10-50s.
		† Jubæa spectabilis	2-10-25s.
		† — <i>Toralyi (very rare)</i>	100s.
		† Kentia australis	10s.



Cocos Weddelliana.

Kentia Balmoreana (<i>Griesebachia</i>)	5-10-25s.	† Kentia divaricata (Illustr.).	25-50s.
†—Canterburyana (<i>Hedyscape</i>) 10-25s.			



Kentia divaricata.



Kentia Lindeni.

† <i>Kentia Forsteriana</i> (<i>Griesbachia</i>)	5-10-25s.	† <i>Latania Commersoni</i>	15-15s.
— <i>gracilis</i> (<i>Illustr.</i>)	25-50s.	— <i>glaucophylla</i>	10-25s.
† — <i>Lindeni</i> (<i>Illustr.</i>)	30-100s.	— <i>rubra</i>	10s.
† — <i>Lucianii</i> (<i>Illustr.</i>)	50-100s.	<i>Licuala elegans</i>	10s.
† — <i>Mac Arthuri</i>	20-50s.	— <i>grandis</i> (<i>Pritchardia</i>) (<i>Illustr.</i>)	125s
† — <i>Mooreana</i>	25-50s.	— <i>peltata</i>	10s.
† — <i>rupicola</i>	10-50s.	<i>Livistona altissima</i>	10-25s.
† — <i>robusta</i> (<i>Cyphokenzia</i>)	10-25s.	† — <i>australis</i> (<i>Corypha</i>)	1-3-5s.
† — sp. <i>Sydneyi</i>	30s.	— <i>Hoogendorpi</i>	1-5s.
<i>Korthalsia</i> (<i>Calamosagus</i>)		† — <i>Jenkinsiana</i>	25s.
— <i>Junghuhni</i>	10-25s.	† — <i>mauritiana</i>	10s.
<i>Latania aurea</i> (<i>Illustr.</i>)	15-25s.	† — <i>olivæformis</i>	3-10s.
† — <i>borbonica</i>	1-5s.	— <i>rotundifolia</i>	10-25s.
		† — <i>sinensis</i> (<i>Latania borbonica</i>)	1-5s.



Kentia robusta.

Livistona subglobosa . . . 15-25s.

Martinezia (Aiphanes)

— *caryotaefolia* 10s.

† — *disticha* 5-10s.

— *erosa* 10-25s.

Maximiliana regia (Illustr.) 5-25s.

Metroxylon elatum 18s.

Oreodoxa (Gaussia) Ghies-

breghti (*O. ventricosa*) . . . 25s.

— *regia* (Illustr.) 2-5s.

— *Sanchona (Euterpe edulis)* 1-5s.

— — *vera* 10s.

Phœnicophorium Sechel-

larum (Stevensonia

grandifolia) (III.) . . . 10-15s.


— — *viridifolium* . . . 15-50s.

Phœnix Andersoni (new.) 100s.




Phœnix reclinata.

+ <i>Phœnix dactylifera</i>	2-10s.	<i>Pinanga maculata</i> (Illustr.)	10s.
— <i>leonensis</i>	5-10s.	— <i>rubricaulis</i>	30-50s.
+ — <i>reclinata vera</i>	1-3-5s.	— <i>sumatrana</i>	50s.
+ — <i>rupicola</i> (Illustr.)	10-25s.	— <i>ternatensis</i>	25s.
— <i>senegalensis</i>	3s.	— <i>Veitchi</i>	25s.
— <i>tenuis</i>	10s.	<i>Plectocomia assamica</i>	50s.
<i>Phytelephas macrocarpa</i>	25-150s.	<i>Pritchardia aurea</i>	15-50s.
<i>Pinanga caesia</i>	75s.	+ — <i>filifera</i> (Illustr.)	2-3s.
— <i>Kuhlii</i> (<i>Ptychosperma</i>)	10-25s.		



Phœnix rupicola.



Licuala (Pritchardia) grandis.

<i>Pritchardia (Licuala) grandis</i>	125s.
— <i>Gaudichaudi</i>	30-150s.
— <i>macrocarpa</i>	15-150s.



Pritchardia macrocarpa.

<i>Pritchardia Martiana</i>	15-50s.	<i>Sabal neocine</i>	2s.
— <i>pacifica</i>	15-25s.	† — <i>princeps</i>	3-10s.
<i>Ptychosperma Alexandræ</i>	1-5s.	— <i>Sanfordi</i>	5s.
— <i>rupicola</i>	30-50s.	— <i>umbraculifera</i>	3-10s.
<i>Ravenea Hildebrandti</i> (III.)	40-100s.	† <i>Seaforthia elegans</i> (III.)	2-5s.
† <i>Rhapis flabelliformis</i>	3-10s.	† — <i>robusta</i> (<i>Areca Baueri</i>)	2-15s.
† — — <i>foliis variegatis</i>	50s.	<i>Synechanthus fibrosus</i>	25s.
† — — <i>cochininchinensis</i>	10s.	<i>Thrinax barbadensis</i> (III.)	50s.
† <i>Sabal Adansoni</i>	2s.	† — <i>Chuco</i>	15-75s.
† — <i>Blackburniana</i>	2-5s.	† — <i>elegans</i>	2-10s.
— <i>cœrulescens</i>	15-25s.	— <i>elegantissima</i>	10-25s.
† — <i>dealbata</i>	3s.	— <i>gracilis</i>	10-50s.
— <i>havanensis</i>	2s.	— <i>graminifolia</i>	10s.
— <i>Magdalenaæ</i>	3-10s.	— <i>grandis</i>	30s.
— <i>mauritiæformis</i>	2-5s.	— <i>Martii</i>	10-30s.

<i>Thrinax parviflora</i>	15-50s.	<i>Verschaffeltia splendida</i>
— <i>radiata</i>	15s.	(Illustr.) 5-15s.
+ <i>Trithrinax aculeata</i> (<i>Chamaerops stauracantha</i>) . .	10-25s.	<i>Wallichia caryotoides</i> 25s.
+ — <i>brasiliensis</i> (<i>Thrinax Chuco</i>)	15-75s.	— <i>myriostigma</i> 50s.
<i>Verschaffeltia melanochætes</i> (<i>Roscheria</i>) (III.) . .	3-25s.	— <i>oblongifolia</i> 40s.
		— <i>porphyrocarpa</i> 20s.
		<i>Welfia regia</i> (<i>Wendl.</i>) (III.) . . 25-50s.
		<i>Zalacca edulis</i> fort. 25s.

We possess the largest collection of Palms in Europe; Gentlemen who are desirous of forming a collection of specimens for show purposes will do well to communicate with us, when we will send list with dimensions, prices and varieties.

PALMS IN STORE POTS.

Large varieties of all the best kinds.



COLLECTION OF PALMS

1st Collection. — Choice Palms.

- | | |
|--------------------------------------|---|
| 1. <i>Pritchardia pacifica</i> . | 7. <i>Latania aurea</i> . |
| 2. <i>Areca nobilis</i> . | 8. <i>Astrocaryum Mexicanum</i> . |
| 3. <i>Kentia Lindeni</i> . | 9. <i>Latania rubra vera</i> . |
| 4. <i>Pinanga maculata</i> . | 10. <i>Geonoma Ghiesbreghtii</i> . |
| 5. <i>Dæmonorops melanochætes</i> . | 11. <i>Pinanga Kuhlei</i> . |
| 6. <i>Verschaffeltia splendida</i> . | 12. <i>Phœnicophorium Sechellarum</i> . |

Price : 80s.

This same collection is available in larger specimens for 120s.

2nd Collection.

- | | |
|---------------------------------|--|
| 1. <i>Arenga saccharifera</i> . | 7. <i>Chamædorea Ernesti-Augusti</i> . |
| 2. <i>Sabal Sandfordi</i> . | 8. " lunata. |
| 3. <i>Kentia Balmoreana</i> . | 9. <i>Oreodoxa Sanchona vera</i> . |
| 4. <i>Cocos Romanzoffiana</i> . | 10. " regia. |
| 5. <i>Phœnix rupicola</i> . | 11. <i>Verschaffeltia melanochætes</i> . |
| 6. <i>Calamus marginatus</i> . | 12. <i>Wallichia porphyrocarpa</i> . |

Price : 50s. and 80s.

3rd Collection.

- | | |
|---------------------------------|------------------------------------|
| 1. Kentia robusta. | 4. Morenia Lindeni. |
| 2. Korthalsia Junghuhni. | 5. Dæmonorops hystrix. |
| 3. Thrinax Chuco. | 6. Drymophlœus olivæformis. |

Price : 35s. and 60s.

4th Collection.

Comprising varieties suitable for apartments.

- | | |
|-------------------------------|-----------------------------------|
| 1. Latania borbonica. | 5. Areca lutescens. |
| 2. Geonoma gracilis. | 6. Phœnix reclinata. |
| 3. Corypha australis. | 7. Cocos flexuosa. |
| 4. Seaforthia elegans. | 8. Ptychosperma Alexandræ. |

In fine plants about 15 inches high : 15s. In stronger specimens 30s. and 50s.

5th Collection.

Comprising very choice varieties for amateurs.

- | | |
|----------------------------|------------------------------|
| 1. Licuala grandis. | 4. Weitchia Johannis. |
| 2. Jubaea Torally. | 5. Thrinax Chuco. |
| 3. Calamus Lindeni. | |

The collection of six varieties : 300s.

The General Catalogue containing a more extensive selection of Palms will be sent on application being made.

**STOVE PLANTS.**

SELECTION of the best Stove and temperate house plants.

The Establishment possesses a large collection of specimens and half-specimens of *Stove-Plants*.

Prices on application.

Those plants which are of recent introduction are marked with an *.

Abutilon <i>Sellowianum marmo-</i>	*Anthurium <i>Schertzerianum</i>
ratum 1-25s.	var. <i>maximum</i> 150s.
Acalypha <i>macrophylla</i> (III.) 2-5s.	*— — — <i>pygmaeum</i> 15s.
— <i>musaica</i> 2-25s.	*— — — <i>Rothschildianum</i> 100s.
— <i>tricolor</i> 2-25s.	*— — — — <i>de la Devan-</i>
Adelaster <i>albo-venosa</i>	<i>sayanum</i> 250-500s.
Adelobotrys <i>Lindeni</i> 3-5s.	*— — — <i>Williamsi</i> (III.) 5-50s.
* Aglaonema (<i>Schismatoglottis</i>) <i>La Valleei</i> (Illustr.) 10-25s.	*— — <i>trilobum</i> (Illustr.) 15-25s.
*— <i>pictum</i> (Illustr.) 15s.	*— <i>Veitchii</i> (Illustr.) 50s.
*— <i>Rebellini</i> 10-25s.	*— <i>Warocquei</i> (Illustr.) 10-25s.
Allamanda <i>Chelsoni</i> 5s.	Aphelandra <i>aurantiaca</i> 2s.
— <i>grandiflora</i> 5s.	— <i>Roezlii</i> 1-5s.
— <i>Hendersoni</i> (Illustr.) 5s.	— <i>fascinata</i> (Illustr.) 2-15s.
Alocasia <i>Boryana</i> 3-25s.	— <i>Leopoldi</i> 1-10s.
— <i>gigantea</i> 3-25s.	— <i>nitens</i> 5s.
— <i>hybrida</i> 15s.	*— <i>punctata</i> 5s.
*— <i>Johnstoni</i> (Illustr.) 10-25s.	— <i>Sinitzini</i> (Illustr.) 2-10s.
— <i>Lowi</i> 5-20s.	* Aralia <i>Chabrierii</i> 10-50s.
— <i>macrorhiza variegata</i> (III.) 3-25s.	*— <i>cochleata</i> 15-25s.
— <i>Marshalli</i> 5-15s.	— <i>elegantissima</i> (Illustr.) 4-75s.
— <i>metallica</i> (Illustr.) 3-20s.	— <i>filicifolia</i> (Illustr.) 3-50s.
*— <i>Putzeysi</i> (Illustr.) 75-200s.	— <i>gracillima</i> (Illustr.) 5-50s.
— <i>Sedeni</i> (Illustr.) 3-10s.	— <i>Guilfoylei</i> 2-25s.
*— <i>singaporensis</i> 10-50s.	*— <i>Massangeana</i> 10s.
*— <i>Thibauti</i> (Illustr.) 15-25s.	*— <i>monstrosa</i> 15s.
*— <i>Veitchi</i> (Illustr.) 5-25s.	— <i>Nilsoni</i> 25-50s.
*— <i>zebrina</i> 5-25s.	— <i>nobilis</i> 25s
Alloplectus <i>Schlimi</i> 3-25s.	*— <i>queratifolia</i> 10-25s.
* Amorphophalus <i>Lacourii</i> (III.)	*— <i>reginæ</i> (Illustr.) 50s.
	— <i>reticulata</i> 1-10s.
— <i>Rivieri</i> 5s.	— <i>sonchifolia</i> (Illustr.) 25s.
* Anthurium <i>Andreanum</i> (Illustr.) 15-250s.	— <i>spectabilis</i> (Illustr.) 50s.
— <i>crystallinum</i> (Illustr.) 5-250s.	— <i>Veitchi</i> (Illustr.) 3-25s.
— <i>Dechartrei</i> (<i>Spathiphyllum canaeefolium</i>) (Illustr.) 2-50s.	Ardisia <i>crenulata</i> 1-5s.
— <i>floribundum</i> (<i>Spathiphyllum</i>) (Illustr.) 5-10s.	— <i>metallica</i> (Illustr.) 5s.
*— <i>insigne</i> 25s.	Aristolochia <i>clypeata</i> (III.) 10s.
— <i>leuconeurm</i> (Illustr.) 3-25s.	— <i>Duchartrei</i> (Illustr.) 5s.
*— <i>Lindenianum</i> (Illustr.) 25-50s.	Artanthe <i>magnifica</i> (III.) 2-15s.
— <i>magnificum</i> 5-25s.	Artocarpus <i>Cannonii</i> (III.) 2-20s.
— <i>ornatum</i> 10-50s.	— <i>grandis</i> 10-50s.
— <i>regale</i> 5-25s.	Asparagus <i>plumosus</i> 5s.
— <i>Schertzerianum</i> 3-25s.	*— <i>nanus</i> 25s.
— — <i>var. Adriani</i> (Illustr.) "	— <i>racemosus</i> 5s.
*— — — <i>grandiflorum</i> 15-50s.	— <i>tenuifolius</i> 25-50s.
*— — — <i>giganteum</i> 50-75s.	Ataccia <i>cristata</i> 10-25s.
	* Batatas <i>paniculata</i> 5-10s.
	* Begonia <i>amabilis</i> 10s.
	*— <i>diadema</i> 3-5s.
	*— <i>Meysseliana</i> 5s.

*— <i>Begonia</i> Teuscheri	4s.	<i>Centropogon</i> Luciani	1-10s.
*— — <i>var. punctata</i>	4s.	<i>Centrosolenia</i> bullata (III.) .	3-5s.
*— — — <i>maculata</i>	4s.	<i>Chirita</i> lilacina (Illustr.) .	3-5s.
<i>Bertolonia Disraeli</i>	10s.	— sinensis.	2s.
— <i>formosa</i>	10s.	* <i>Cienkowskia Kirkii</i> (III.) .	25s.
— <i>Gladstonei</i>	5s.	<i>Cissus</i> albo-nitens	2s.
— <i>guttata</i>	5s.	— <i>amazonica</i>	2s.
— — <i>albo punctatissima</i> . .	3s.	— <i>argentea</i>	2s.
— — <i>roseo punctatissima</i> . .	3s.	— <i>argyrea</i>	2s.
— — <i>splendens</i>	20s.	— <i>discolor</i>	1s.
— <i>Killischi</i>	5s.	— <i>Endresi</i>	2s.
— <i>Legrelleana</i>	25s.	— <i>Lindeni</i> (Illustr.)	3s.
— <i>Marchandi</i>	3s.	— <i>porphyrophylla</i>	2s.
— <i>maculata</i>	3s.	* <i>Clausenia corymbiflora</i> . .	10-25s.
— <i>margaritacea</i>	3s.	<i>Clavija ornata</i>	5-50s.
— <i>marmorata</i>	2s.	<i>Clerodendron Balfouri</i>	2s.
— <i>Mirandei</i>	10s.	— <i>fallax</i>	3s.
— <i>Ohlendorffii</i>	3s.	— <i>fragrans flore pleno</i> . .	3s.
— <i>ornata</i>	10s.	— <i>infortunatum</i>	3s.
— <i>pubescens</i>	5s.	<i>Clidemia vittata</i> (Illustr.) .	3-5s.
— <i>Rodecki</i>	5s.	<i>Clivea miniata</i>	2-50s.
— <i>smaragdina</i>	5s.	*— — <i>var. Lindeni</i> (III.) . .	20-50s.
— <i>splendida</i> (Illustr.)	10s.	<i>Cochliostema Jacobianum</i>	
— <i>velutina</i>	5s.	(Illustr.)	5-30s.
— <i>Van Houttei</i>	3s.	<i>Coccoloba macrophylla</i> . . .	10s.
<i>Bignonia aurantiaca</i>	10-15s.	<i>Coffea arabica</i> fol. var. . . .	3-5s.
— <i>argyrea violescens</i> (III.) .	3s.	<i>Colocasia</i> albo-violacea . . .	3s.
— <i>magnifica</i>	15-20s.	— <i>argyroneura</i>	5s.
— <i>ornata</i>	5s.	— <i>macrorhiza</i> fol. var. (III.) .	3-25s.
— <i>picta</i> (Illustr.)	5s.	*— <i>Neo-Guineensis</i> (Illustr.) .	5-50s.
— <i>speciosa</i>	2s.	<i>Coutarea Schereffiana</i> (Illustr.)	
<i>Bombax fastuosa</i>	10s.	hort. 1878)	5-10s.
— <i>insigne</i>	4s.	<i>Cossignya borbonica</i>	10-25s.
<i>Bougainvillea glabra</i>	3s.	<i>Cupanea filicifolia</i>	15-25s.
<i>Brexia chrysophylla</i>	3s.	<i>Curatella (Theophrasta) im-</i>	
<i>Caliphurria subedentata</i> (III.)	3s.	perialis	10-50s.
<i>Callicarpa purpurea</i> (Illustr.)	2-5s.	<i>Curculigo recurvata</i>	1s.
— <i>sumatrana</i>	5s.	— — <i>variegata</i>	5-50s.
<i>Campsodium filicifolium</i> . .	3-5s.	* <i>Cureuma Leopoldi</i>	25s.
<i>Campylobotys Ghiesbreghtii</i>		<i>Curmeria picturata</i> (Illustr.)	3-25s.
(Illustr.)	2-5s.	* — <i>Reezi</i>	5s.
— — <i>fol. var.</i>	2s.	* — <i>Wallisi</i> (Illustr.) . . .	5s.
— — <i>marmorata</i>	2-5s.	<i>Cyanophyllum magnificum</i> .	3-75s.
— — <i>Ortgiesi</i>	3-5s.	— <i>spectandrum</i>	5s.
— — <i>pyrophylla</i>	2-5s.	<i>Cyphonandra argentea</i> . .	3-25s.
— — <i>refulgens</i>	2-5s.	<i>Cyrtodeira cupreata</i>	1-5s.
— — <i>regalis</i>	2-5s.	— <i>fulgida</i> (Illustr.)	1-5s.
* <i>Carissa grandis</i>	10s.	— <i>metalica</i>	1-5s.
<i>Centradenia rosea</i>	3-15s.	<i>Dichorisandra musaica</i> . .	5-25s.



Heliconia aurea striata.

Dichorisandra metallica picta	5-15s.	Fittonia gigantea (Illustr.)	1-5s.
— undata	3-25s.	— Pearcei	1-5s.
— vittata	2-10s.	— Verschaffelti	1-5s.
* Dieffenbachia amoenia	5-15s.	Franciscea Lindeni	3s.
— antioquensis (Illustr.)	10-50s.	Galipea odoratissima	3s.
— Baraqimiana (Illustr.)	3-15s.	Gardenia amoena.	3-5s.
— Bausei (Illustr.)	3-25s.	— citriodora	3s.
— Bowmanni	3-20s.	— florida	3s.
— brasiliensis	10-25s.	— fortuneana	3-5s.
— eburnea	5s.	— globosa	10s.
— gigantea (Illustr.)	3-30s.	— radicans	1-2s.
— illustris (Illustr.)	5s.	— — fol. varg.	3s.
— imperialis (Illustr.)	10-30s.	— Stanleyana	3s.
— lancifolia (Illustr.)	3-10s.	Geissois racemosa (Illustr.)	10-25s.
— latimaculata (Illustr.)	3-50s.	Gloneria jasminiflora (Illustr.)	5-10s.
*— Leopoldi (Illustr.)	5-50s.	* Haemanthus Kalybreyeri (III.)	10s.
*— magnifica (III.)	30-75s.	* Heliconia aurea-striata (III.)	15-50s.
— memoria Corti	15-30s.	— metallica	15s.
— nobilis	5s.	Hibiscus rosa sinensis	3s.
— Parlatorei (Illustr.)	5-50s.	— — — Cooperi (Illustr.)	2-5s.
— — marmorea	10-25s.	— — — flore pleno	2-5s.
— picta	3-50s.	*— — — Lucien Linden (III.)	10s.
— Wallisi (Illustr.)	15-25s.	*— — — schizophetalus	3s.
— Weiri superba	5s.	Hoya bella.	3s.
Dillenia speciosa.	5s.	— carnosa.	2s.
Dioscorea discolor	3s.	— — — fol. var.	3s.
— metallica (Illustr.)	5s.	— imperialis	3s.
Dorstenia argentea	3-10s.	— Paxtoni.	2s.
Drymonia turialvae (Illustr.)	2-15s.	Ixora crocata-rutilans	2-5s.
Echites argyrea (Illustr.)	2-10s.	— odorata.	5s.
— refulgens	3-15s.	Jacaranda mimosaeifolia	7-15s.
— rubro-venia (Illustr.)	2-10s.	* Jasminum gracillimum	5-10s.
* Epipremnum mirabile (Tonga)	10-25s.	— hirsutum	5s.
Eranthemum Andersoni	3s.	Jatropha multifida	15-25s.
— albo-marginatum	5s.	* Kaempferia Gilbertii.	10-15s.
— atropurpureum	2-5s.	Lasiandra floribunda	3s.
— aureo-reticulatum	1-5s.	— macrantha (Illustr.)	2-10s.
— igneum	1-5s.	* Lathyrus splendens	15s.
— nigrum (Illustr.)	3-5s.	Laurus querciformis	10s.
— roseum (Illustr.)	2-10s.	Ledenbergia roseo-aenea (III.)	3-10s.
— sanguineum	2-10s.	Lindenia rivalis (Illustr.)	5s.
— Schomburgki (Illustr.)	1-5s.	Makoya bella (Illustr.)	3-5s.
— tricolor	2-10s.	Medinilla amabilis	10s.
— variabile.	2-10s.	— erythrophylla	10-25s.
Eucharis amazonica (Illustr.)	2-5s.	— magnifica	4-10s.
— candida	3-10s.	Meyenia alba	10s.
Eugenia magnifica	10s.	— erecta (Illustr.)	5s.
Euphorbia jacquiniaeifolia . . .	2-3s.	Menima turgida	10s.
— splendens	3s.	Mimosa pudica (Sensitive)	2s.
* Euryceles australasiaica	3-5s.	Monstera delicosa	5s.
Ficus Anneli	5-10s.	Musa Cavendishi.	10s.
— cerasiformis (Illustr.)	3s.	— coccinea	15-25s.
— dcalbata (Illustr.)	5-15s.	— ensete	2-150s.
— edulis.	5s.	*— sumatrana (Illustr.)	25s.
— elastica	2-5s.	— superba.	10-15s.
— elegans	3s.	— textilis	25s.
— Parcelli	1-5s.	— vittata	50-150s.
— Portecana	3-10s.	— zebrina	25s.
Fittonia argyroneura	1-10s.	Mussaenda frondosa	5s.
		Panax fruticosum	3-5s.

Panax laciniatus	3-5s.	* Schismatoglottis (<i>Aglaonema</i>)	25s.
— albo spicata	5-10s.	<i>crispata</i>	25s.
Pandanus graminifolius . . .	5s.	*— <i>Lavalleana</i> (<i>III.</i>)	10-25s.
— <i>Javanicus</i>	10s.	*— <i>Rebellini</i>	10-25s.
— variegatus	15s.	Sciadocalyx <i>Luciani</i> (<i>III.</i>)	2-10s.
— <i>ornatus</i> (<i>Illustr.</i>)	15s.	Scindapsus <i>pictus</i>	2-5s.
— <i>Panzeri</i> (<i>Illustr.</i>)	10-75s.	Smilax <i>marmorea</i>	5s.
— <i>utilis</i>	1-15s.	<i>vaginata</i>	5s.
— <i>Vandermeerschi</i>	15s.	Sonerila <i>argentea</i>	2s.
— <i>Veitchi</i>	5-200s.	— <i>Alp.</i> <i>Van de Sande</i> (<i>III.</i>)	3s.
Paullinia quassiaeformis . . .	3-5s.	— <i>Baronne de Marche</i> (<i>III.</i>)	3s.
— <i>thalictrifolia</i>	1-3s.	— <i>Hendersoni</i>	2s.
— — — <i>argentea</i>	3s.	— <i>marmorata</i>	2s.
Pavetta borbonica	10-25s.	— <i>Mme Gloner</i>	3s.
— <i>picta</i>	10-25s.	— <i>Mme Ed. Otlet</i> (<i>Illustr.</i>)	3s.
* Pavonia Makoyana	4s.	— <i>Mme Vict. Alesch</i> (<i>III.</i>)	3s.
Pellionia Daveauana (<i>Illustr.</i>)	3s.	— <i>M. Alp. Mame</i> (<i>Illustr.</i>)	3s.
Peperomia argentea	2s.	— <i>margaritacea</i> (<i>Illustr.</i>)	4s.
— <i>argyrea</i>	2s.	— — <i>superba</i>	4s.
— <i>eburnea</i>	2s.	— <i>pictureata</i>	5s.
— <i>maculosa</i>	5s.	*— <i>Lili</i>	4s.
— <i>prostrata</i>	3s.	*— <i>Dora</i>	4s.
— <i>resedaeiflora</i> (<i>Illustr.</i>) . . .	2s.	*— <i>Rouly</i>	4s.
— <i>v. lutina</i> (<i>Illustr.</i>)	2s.	*— <i>Rita</i>	4s.
— <i>Verschaffeltii</i> (<i>Illustr.</i>)	1-3s.	*— <i>Ada</i>	4s.
Philodendron daguenense (<i>III.</i>)	5-75s.	*— <i>Nelly</i>	4s.
— <i>gloriosum</i> (<i>Illustr.</i>)	5-25s.	Spathiphyllum <i>cannaeformium</i> (<i>Dechardi</i>) (<i>III.</i>)	3-50s.
*— <i>Lindeni</i>	10-50s.	*— <i>hybridum</i> (Voir page 6) (<i>Illustr.</i>)	25-150s.
— <i>melanochrysum</i> (<i>Illustr.</i>)	5s.	— <i>Patini</i> (<i>Illustr.</i>)	5-25s.
— <i>pertusum</i>	2s.	*— <i>pictum</i>	3-4s.
Phyllanthus atropurpureus . . .	2s.	Sphaerogyne <i>cinnamomea</i> (<i>III.</i>)	5-15s.
— <i>nivosus</i> (<i>Illustr.</i>)	3s.	— <i>imperialis</i> (<i>Illustr.</i>)	5-25s.
— <i>roseo-pictus</i>	5s.	— <i>latifolia</i>	10-25s.
Phyllogatis rotundifolia	5s.	S'ephanotis <i>floribunda</i>	2-15s.
* Phyllocladus <i>Lindeni</i> (<i>III.</i>)	3-10s.	Strelitzia <i>Nicolai</i>	20s.
* Piper metallicum	10s.	— <i>regina</i>	15s.
Plumbago rosea var. <i>coccinea</i> .	10s.	* Tacsonia <i>Buchananii</i> (<i>Illustr.</i>)	3-5s.
Pogostemon Patchouli	1s.	Theophrasta <i>imperialis</i> (<i>III.</i>)	15s.
Pothos argyreia (<i>Illustr.</i>)	2s.	— <i>Jussieu</i>	15s.
*— <i>aurea</i> (<i>Illustr.</i>)	2-5s.	Tradescantia <i>discolor</i>	1-5s.
*— <i>celatocalyx</i>	5-15s.	— <i>lineata</i>	2s.
Ravenalia madagascariensis (<i>Illustr.</i>)	4-10s.	— <i>Madame Lequesne</i>	1-5s.
Rogiera gratissima	5-10s.	— <i>odoratissima</i>	10s.
— <i>cordata</i>	5s.	— <i>zebrina</i>	2-5s.
Rondeletia major	3s.	Vanilla <i>aromatica</i>	5-25s.
Sanchezia nobilis (<i>Illustr.</i>) . . .	2s.		
— <i>glaucophylla</i> (<i>Illustr.</i>) . . .	2s.		

The General Catalogue containing a more detailed list of Stove plants will be forwarded on application being made.

NEPENTHES.

<i>Nepenthes albo-marginata</i>	25s.	<i>Nepenthes laevis</i>	5-50s.
— <i>ampullacea</i> (<i>Illustr.</i>)	10-30s.	— <i>lanata</i> (Veitchi) price on demand.	
— — <i>vittata major</i> (<i>Ill.</i>)	15-30s.	— <i>Lawrenciana</i> (<i>Illustr.</i>)	50s.
— <i>Anerleyensis</i>	50s.	— <i>maculata</i>	20-30s.
— <i>atro sanguinea</i>	75s.	— <i>madagascariensis</i>	100s.
— <i>bicalcarata</i> (<i>III.</i>)	50-75-125s.	— <i>Morganiae</i>	75s.
— <i>Chelsoni</i>	25-50s.	— <i>Outramiana</i>	50s.
— <i>coccinea</i>	75s.	— <i>phyllamphora</i>	10-25s.
— <i>compacta</i>	50s.	— <i>Rafflesiana</i>	15-50s.
— <i>Courtii</i>	25s.	— <i>robusta</i>	50s.
— <i>distillatoria</i> (Kharysiana)	5-10s.	— <i>rubra</i>	25-50s.
— <i>Dominiana</i>	10-50s.	— <i>Rajah</i>	150s.
— <i>Dormaniana</i>	80s.	— <i>Ratcliffiana</i>	75s.
— <i>gracilis-major</i>	5-10s.	— <i>rubro-maculata</i>	50s.
— <i>Harryana</i>	50s.	— <i>sanguinea</i> , price on demand.	
— <i>Henryana</i> (<i>Illustr.</i>)	50s.	— <i>Sedenii</i>	5-20s.
— <i>Hookerae</i>	25s.	— <i>rubra</i>	25-50s.
— <i>Hookeriana</i>	15-25s.	— <i>Stewartii</i>	30s.
— — <i>elongata</i>	25s.	— <i>superba</i>	50s.
— <i>hybrida</i>	15s.	— <i>Veitchii</i> (<i>lanata</i>) price on demand.	
— — <i>maculata</i>	15s.	— <i>Williamsi</i>	50s.
— <i>intermedia</i>	15-20s.	— <i>Wrigleyana</i>	75s.
— <i>Kennedyana</i>	25s.	— <i>zeylanica-rubra</i> (<i>hirsuta</i>)	30s.

Collection of 6 nice plants, 40, 75, 150 shillings.

Collection of 12 nice plants with pitchers 175, 300 shillings.

Specimens furnished at low prices for Exhibition purposes.



ORCHIDS.



We give here a selection of the best varieties only, for further varieties see our special catalogue of Orchids which will be sent on application being made.

Our collection of these lovely plants is the largest in Europe and every day we are increasing the stock, special attention is paid to rare and unique varieties.

How often have we heard said that Orchids are such difficult plants to cultivate, whereas with a few rare exceptions they are among the easiest of all house plants, and then where will one find another class of plants so graceful, so sweetly perfumed and so curious? Of all plants they are the

best for decorating rooms when in flower; some, among which we would especially recommend the Odontoglossums, Oncidioms and Dendrobiums, stay for months in flower thus well repaying any slight trouble attendant on their culture.

They will thrive in any house, whether a simple little viney or the colossal greenhouse. Suspended and intermingled among ferns, palms and other plants they brighten up with their gorgeous blooms what would otherwise be but a dull mass of green.

To the Botanist what plants afford more interest and instruction in studying the marvellous contrivances for their fertilisation and propagation.

For the benefit of those who wish to become amateurs of this lovely class of plants and who are hindered by not knowing what varieties to chose we have formed a series of cheap collections which they will find easy to cultivate and yet consisting of some of the most lovely forms to be found.

CHOICE OF THE BEST VARIETIES OF

ORCHIDS

N. B. Our special catalogue will be sent on application being made.

COOL ORCHIDS.

Those cultivated in a hot temperature are marked with a (c).

Those cultivated in a temperate atmosphere are marked with a (t).

The varieties introduced by the firm are marked with an (*).

*Aclineta , Ldl.		Cymbidium , Swartz.
— Barkeri	15s.	— affine. 125s.
Acropera , Ldl.		— aloefolium 10-25s.
— Loddigesii	10s.	— ensifolium 20s.
*Ada , Ldl.		— grandiflorum 10-25s.
— aurantiaca (Illustr.) . . .	10-20s.	— Hookerianum 40-75s.
*Anguloa , R. et P.		— longifolium. 30-50s.
— Clowesi (Pescat)	15-50s.	— Parishi 40-100s.
*— Ruckeri	25-40s.	— pendulum 10s.
*— (<i>purpurea</i>) sanguinea (III.)	75s.	— sinense 10s.
*— uniflora	25-50s.	 Cypripedium , Linn.
— — superba	30s.	— barbatum 5-10s.
*Arpophyllum , La Llav.		— — atropurpureum 25s.
— cardinalis (Pescat) . . .	50s.	— — grandiflorum 10-25s.
— giganteum	25-100s.	 Disa , Lin.
*— spicatum	15s.	— grandiflora (Pescat) . . . 10-25s.
Barkeria , Know. et West.		 Epidendrum , Lin.
— elegans (Illustr.) (Pescat)	25s.	*— Frederici-Guilhelmi (III.) . . . 75s.
— spectabilis	15-25s.	— glumaceum 50s.
Cattleya , Ldl.		— leucochilum 50s.
— citrina	15s.	*— paniculatum 20s.
		— — splendens 50-100s.

* <i>Epidendrum pentotes</i>	25s.	<i>Masdevallia infracta</i>	10-25s.
— prismatocarpum	20-50s.	*— <i>Lindeni</i> (<i>Illustr.</i>)	10-25s.
— <i>pseudo-Epidendrum</i>	100s.	*— <i>maculata</i>	10-25s.
— <i>radiatum</i>	25s.	*— <i>macrura</i>	150s.
*— <i>Ruckeri</i>	25s.	*— <i>melanopoda</i>	10-20s.
*— <i>sceptrum</i>	20-75s.	*— <i>nycterina</i> (<i>Illustr.</i>)	15s.
*— <i>Sophronitis</i>	30-50s.	*— <i>ochtodes</i>	10-25s.
— <i>vitellinum</i> (<i>Illustr.</i>)	10-30s.	*— <i>Peristeria</i> (<i>Illustr.</i>)	20s.
— — <i>majus</i>	15-75s.	*— <i>polysticta</i> (<i>Illustr.</i>)	10-40s.
<i>Evelyna</i> , Ldl.		*— <i>Tovarensis</i> (<i>Cand.</i>) (<i>III.</i>)	10-50s.
*— <i>kermesina</i>	50s.	— <i>triangularis</i>	25s.
<i>Helcia</i> , Ldl.		*— <i>Trochilus</i> (<i>Colibri</i>) (<i>III.</i>)	50-100s.
*— <i>sanguinolenta</i> (<i>Illustr.</i>)	1-30s.	— <i>troglodytes</i>	35-50s.
<i>Isochilus graminifolius</i>	25s.	— <i>Veitchiana</i>	15-50s.
<i>Lælia</i> , Ldl.		Maxillaria , R. et Pav.	
— <i>acuminata</i>	15s.	*— <i>grandiflora</i> (<i>Illustr.</i>)	15-25s.
— <i>albida</i>	10-30s.	— <i>Harrisoniana</i>	10-25s.
— <i>anceps</i>	10-25s.	— <i>luteo-alba</i>	15s.
— <i>autumnalis</i> (<i>Illustr.</i>)	10-30s.	*— <i>nigrescens</i>	10s.
— <i>cinnabarina</i>	10-25s.	— <i>picta</i>	10-30s.
— <i>flava</i>	15s.	— <i>splendens</i>	50-75s.
— <i>furfuracea</i>	15s.	— <i>tenuifolia</i>	10s.
— <i>majalis</i>	10s.	*— <i>venusta</i> (<i>Pescat.</i>)	15s.
*— <i>superbiens</i>	25-30s.	Mesospinidium , Rehb.	
<i>Lycaste</i> , Ldl.		— <i>sanguineum</i>	10s.
— <i>aromatica</i>	10-20s.	*— <i>vulcanicum</i>	20s.
*— <i>costata</i>	30s.	Odontoglossum , H. B. et Kth.	
— <i>cruenta</i>	10s.	— <i>Alexandriæ</i> (<i>Crispum</i>)	15-50s.
— <i>D'ppei</i>	10s.	— <i>Andersonianum</i>	150s.
— <i>flavescens</i>	15s.	*— <i>augustatum</i>	25s.
*— <i>gigantea</i>	15-30s.	*— <i>aureo-purpureum</i>	25s.
*— <i>Harrisoniana</i>	10-20s.	— <i>bicornutense</i>	10-30s.
*— <i>lanipes</i>	10-25s.	*— <i>album</i> (<i>Illustr.</i>)	25s.
*— <i>macrobolbon</i>	15s.	*— <i>brevifolium</i> (<i>Illustr.</i>)	50-150s.
*— <i>macrophylla</i>	15s.	— <i>Cervantesi</i> (<i>Illustr.</i>)	10-25s.
*— <i>Skinneri</i> (<i>Pescat.</i>)	10-25s.	— <i>maximum</i> (<i>III.</i>)	50s.
— — <i>alba</i> (<i>Illustr.</i>)		— <i>roseum</i>	25s.
<i>Masdevallia</i> R. et Pav.		*— <i>cirrhosum</i> (<i>Illustr.</i>)	10-30s.
*— <i>amabilis</i>	10-30s.	— <i>Klabochorum</i>	30-60s.
*— <i>var. lineata</i> (<i>Illustr.</i>)	25-100s.	*— <i>citrosum</i> (<i>Illustr.</i>)	10-30s.
— <i>bella</i>	10-100s.	*— <i>constrictum</i>	50s.
*— <i>Benedicti</i>	15-50s.	*— <i>cordatum</i> (<i>Pescat.</i>)	10-25s.
— <i>Brückmulleri</i>	5-10s.	*— <i>coronarium</i> (<i>Pescat.</i>)	50-125s.
*— <i>candida</i> (<i>Tovarens.</i>) (<i>III.</i>)	10-25s.	— <i>miniatum</i>	50-100s.
*— <i>caudata</i>	10-25s.	— <i>crispum</i> (<i>Alexandriæ</i>)	15-50s.
*— <i>chimæra vera</i> (<i>Illustr.</i>)	45-150s.	*— <i>var. Mariæ</i> (<i>Illustr.</i>)	100s.
*— <i>civilis</i>	5-10s.	*— <i>cristatum</i> (<i>Illustr.</i>)	15-50s.
*— <i>coccinea</i>	25-50s.	— <i>crocatum</i>	50s.
— <i>corniculata</i>	40-80s.	*— <i>densiflorum</i>	25-50s.
— <i>cucullata</i>	2-50s.	— <i>Ehrenbergii</i> (<i>Illustr.</i>)	15-30s.
— <i>Davissii</i>	20-50s.	— <i>gloriosum</i>	25-75s.
*— <i>elephanticeps</i>	75-150s.	— <i>grande</i>	10-30s.
*— <i>fenestrata</i>	15s.	*— <i>Halli</i> (<i>Illustr.</i>)	25-50s.
*— <i>Gorgona (nycter)</i> (<i>III.</i>)	15-50s.	*— <i>hastilabium</i> (<i>Pescat.</i>)	20-50s.
— <i>Harryana</i> (<i>Illustr.</i>)	10-75s.	— <i>Insleayi</i>	15s.
*— <i>Calenderi</i>	50-100s.	*— <i>Lindleyanum</i>	10-25s.
*— <i>sanguinea</i>	50-100s.	— <i>speciosum</i>	50s.
— <i>Houtteana</i>	25-100s.	*— <i>Lindeni</i>	40-75s.
*— <i>ignea</i> (<i>Illustr.</i>)	15-50s.	*— <i>Londesburghianum</i> (<i>III.</i>)	15-30s.
*— — <i>Boddaerti</i> (<i>Illustr.</i>)	100s.	*— <i>luteo-purpureum</i> (<i>Illustr.</i>)	20-75s.
		(<i>hystrix</i>)	

*Odontoglossum maculatum (Pescat.)	10s.	Oncidium leucochilum splen-
— Madrense	100s.	dens 30s.
*— nebulosum (Illustr.) . . .	20-75s.	— linguiforme 25s.
*— nevadense (Illustr.) . . .	200s.	macranthum (Illustr.) 30-100s.
*— odoratum (Illustr.) . . .	15-30s.	*— nubigenum (Illustr.) 25s.
*— Pescatorei (Pescat.) . . .	10-50s.	— Batemanianum 30s.
— — immaculatum	50s.	*— Lindleyanum 30s.
— — pulchellum	10-30s.	— ornithorynchum 15s.
— — majus	20-50s.	— Phalaenopsis (Illustr.) 75s.
— — radiatum	50s.	— pretextum 25-50s.
*— Reichenheimi (Pescat.) . . .	30-50s.	— sarcodes (Illustr.) 30-150s.
— Roezli (Illustr.)	15-50-100s.	*— superbiens 50s.
— — album	50-100s.	*— tigrinum (Barkeri) (III.) 10-30s.
*— roseum (Illustr.)	15-25-100s.	— uniflorum 15s.
— Rossi	10-20s.	— varicosum 20-30s.
— — majus (Illustr.)	25-50s.	— — Rogersi 30-50s.
— — superbium	50-100s.	— Weltoni 15-40s.
*— Schlieperianum	20s.	*— zebrinum (Illustr.) 30-80s.
— Schlumi		Pilumna , Ldl.
— spectrum	50-100s.	*— fragrans 15s.
*— triumphans (Illustr.)	20-50s.	*— nobilis (Illustr.) 25-50s.
*— vexillarium (Illustr.)	75-150s.	Pleione , Don.
Oncidium , Sw.		— lagenaria (Illustr.) 10-25s.
*— aurosum (Illustr.)	10-25s.	Restrepia , H. et Kth.
*— Barkeri (tigrin.) (III.) . . .	10-25s.	*— artennifera (Illustr.) 25-100s.
— — concolor (Illustr.)	25s.	— — aurea 25-100s.
— — crispum	10-30s.	*— elegans 25s.
— — grandiflorum	15s.	— — ophiocephala 15s.
— — marginatum	35s.	Sobralia , R. et Pav.
*— cucullatum flavidum	15-30s.	— Cattleya 300s.
*— — maculatum	20-40s.	— macrantha 20s.
— — flexuosum	15-50s.	— — nana
— — hastatum	15-30s.	Sophronitis , Ldl.
— — incurvum (Illustr.)	15s.	— coccinea 20s.
— — — album (Illustr.)	500s.	— — grandiflora (<i>militaris</i>) 15-25s.
— — lepidum	50s.	Thunia , Rchb.
— — leucochilum	15s.	— alba 10s.
		— Bensoniae 10s.

HOTHOUSE AND TEMPERATE ORCHIDS

Those cultivated in a hot temperature are marked with a (c).

Those cultivated in a temperate atmosphere are marked with a (t).

The varieties introduced by the firm are marked with an (*).

Acineta , Lld.		cAerides Fieldingi (<i>Foxbr.</i>) 30-15-50s.
t — superba (<i>Humboldti</i>) (III.) . . .	10-25s.	c — Houlleti (Illustr.) 25-100s.
Aerides , Lour.		*t — japonicum (Illustr.) 15-50s.
c — affine	15-25s.	c — Larpentæ 50-100s.
c — — roseum (Illustr.)	10-25s.	— Leeananum 30-50s.
c — — crassifolium	20-50-100s.	c — Lindleyanum 20-50s.
c — — crispum (Illustr.)	20-50s.	c — Lobbi (Illustr.) 25-125s.

<i>c Aerides maculosum (Pesc.)</i>	25s.	<i>t Cattleya citrina (Pescant.)</i>	10-25s.
<i>c — odoratum (cornutum)</i>	10-25s.	<i>t — crispa (Laelia)</i>	10-25s.
* <i>c — quinquevulnerum</i>	20-75-150s.	* <i>t — dolosa (Illustr.)</i>	25s.
* <i>c — Reichenbachianum</i>		<i>t — Dowiana (Illustr.)</i>	50-100s.
— Schröderi	1500s.	* <i>t — Eldorado (Illustr.)</i>	30-75s.
<i>c — suavissimum</i>	20-50s.	* <i>t — splendens (Illustr.)</i>	50-100s.
<i>c — var. flavidum</i>	50s.	<i>t — elegans (Laelia) (III.)</i>	25-125s.
<i>c — Veitchi</i>	100-300s.	<i>t — Gascaliana (new)</i>	75-150s.
<i>c — virens</i>	15-50s.	* <i>t — gigas</i>	25-50-150s.
<i>c — Dayanum</i>	25-75s.	<i>t — granulosa</i>	30s.
<i>c — var. de Moulmein</i>	50s.	<i>t — guttata</i>	20s.
<i>c — Warneri</i>	50-100s.	<i>t — Harrisoniana</i>	20-50s.
<i>Angræcum</i> , Thouars.		<i>t — violacea</i>	50s.
<i>c — citratum</i>	25s.	<i>t — intermedia</i>	10-25s.
<i>c — eburneum</i>	25-150s.	<i>t — labiata</i>	25-50s.
<i>c — superbum</i>	50-200s.	<i>t — atro-purpurea</i>	50s.
<i>c — Ellisii</i>	50s.	<i>t — Luddemanniana</i>	75s.
<i>c — sesquipedale (Illustr.)</i>	50-200s.	<i>t — — splendens</i>	75s.
<i>c — superbum, strong</i>	150s.	<i>t — — Percivaliana</i>	50s.
<i>Ansellia</i> , Ldl.		<i>t — Leopoldi (Illustr.)</i>	15-50s.
<i>c — africana</i>	25-50s.	<i>t — lobata (Bootheana)</i>	15-50s.
<i>Aspasia</i> , Ldl.		<i>t — Laddemanniana</i>	25s.
<i>t — epidendroides</i>	15s.	<i>t — luteola</i>	25s.
<i>Batemania</i> , Ldl.		<i>t — marginata (Laelia)</i>	20s.
<i>c — meleagris (Illustr.)</i>	30-50s.	* <i>t — maxima (Illustr.)</i>	25-75s.
<i>c — Lobbi</i>	10-15s.	<i>t — Mendeli</i>	20-100s.
<i>Bollea celestis</i>	40-75s.	<i>t — — superba</i>	100s.
— Patini	50s.	* <i>t — Mossiae</i>	10-25s.
<i>Brassavola</i> , R. Br.		<i>t — — Ainsworthi</i>	25-75s.
<i>c — Dibgyana</i>	15-25-100s.	* <i>t — chirguensis</i>	50-100s.
<i>t — glauca</i>	10-25s.	<i>t — — grandiflora</i>	40s.
<i>Brassia</i> , R. Br.		<i>t — — pallida</i>	25s.
* <i>t — caudata Hieroglyphica. (III.)</i>		<i>t — — speciosissima</i>	25-50s.
<i>t — Gireoudiana</i>	25s.	* <i>t — Schlimi</i>	25s.
<i>t — maculata</i>	15s.	* <i>t — superba</i>	30s.
<i>t — verrucosa</i>	10-15s.	— <i>Nobilior (Illustr.)</i>	50s.
<i>Calanthe</i> , R. Br.		<i>t — Perrini (Laelia)</i>	15-30s.
* <i>t — bracteata (new)</i>	50s.	<i>t — Pinelli (Laelia)</i>	15s.
<i>t — Masuca</i>	15-25-50s.	<i>t — pumila (Laelia) (Illustr.)</i>	25-50s.
— — <i>grandiflora</i>	25s.	<i>i — purpurata (Laelia)</i>	15-50s.
<i>t — sylvatica</i>	25s.	<i>t — Brysiniana (Laelia)</i>	50s.
<i>t — Veitchi</i>	15-30s.	<i>t — Leopoldi</i>	75s.
<i>t — — superba</i>	35s.	* <i>t — quadricolor (Illustr.)</i>	25-50s.
<i>t — veratrifolia</i>	15-25s.	* <i>t — carnea</i>	25-40s.
<i>t — vestita</i>	10s.	<i>t — Sanderiana</i>	
<i>t — — luteo-oculata</i>	10s.	<i>t — Schilleriana (Laelia)</i>	30-75s.
<i>Catasetum</i> , L. C. Rich.		<i>t — Skinneri</i>	10-25s.
* <i>c — laminatum</i>	25s.	<i>t — speciosissima (Luddemani-</i>	
* <i>c — Ollare</i>	25s.	<i>niana)</i>	50s.
<i>c — trimerochilum</i>		<i>t — — superba</i>	
* <i>c — Trulla</i>	30s.	* <i>t — — splendida (Illustr.)</i>	30-75s.
<i>Cattleya</i> , Ldl.		* <i>t — — Trianæa</i>	15-25-50s.
<i>t — Acklandiae (Illustr.)</i>	15-50s.	* <i>t — — albida</i>	25-50s.
* <i>t — amethystiglossa</i>	30-75s.	* <i>t — — delicata</i>	25-50s.
(Illustr.)		* <i>t — — virginalis (Illustr.)</i>	50s.
<i>t — amethystina</i>	15-30s.	* <i>t — — Wagneri</i>	75s.
* <i>t — aurea</i>	25-100s.	<i>t — — Walkeriana (bulbosa)</i>	15-50s.
<i>t — bulbosa (Walkeriana)</i>		<i>t — — Wallisii</i>	50-100s.
(Pescat.)	15-50s.	<i>t — — Warneri</i>	
* <i>t — chocensis (Illustr.)</i>	30s.	<i>t — — Warscewiczi</i>	25-75s.
* <i>t — Chocensis Miss Nilsson</i>	50s.		

<i>t</i> <i>Cattleya</i> <i>xanthina</i> (<i>Læla</i>)	25s.	<i>t</i> <i>Cypripedium niveum</i> (III.)	15-30s.
<i>Chysis</i> , Lodd.		<i>t</i> — <i>Parishi</i> (Illustr.)	30-75s.
<i>t</i> — <i>aurea</i>	15-25s.	<i>t</i> — <i>Pearcei</i> (<i>Selenipedium</i>)	15-25s.
<i>t</i> — <i>bractescens</i> (Illustr.)	15-25s.	* <i>t</i> — <i>Roezli</i> (<i>Selenip.</i>) (III.)	20-50s.
<i>Cirrhopetalum</i> , Ldl.		* <i>t</i> — <i>Schlimg</i> (<i>Selenipedium</i>) (<i>Pescat.</i>)	30-75s.
<i>t</i> — <i>elegans</i>	100s.	* <i>t</i> — <i>album</i> (Illustr.)	75-150s.
<i>t</i> — <i>Makoyanum</i>	40s.	<i>t</i> — <i>Sedeni</i>	20-75s.
<i>t</i> — <i>glutinosum</i>	"	<i>t</i> — <i>Spicerianum</i> (Illustr.)	50s.
<i>t</i> — <i>picturatum</i>	25s.	<i>t</i> — <i>Stonei</i> (Illustr.)	50s.
— <i>Thouarsi</i>	25s.	<i>t</i> — <i>superbiens</i> (<i>Veitchianum</i>) (Illustr.)	40-75s.
* <i>t</i> — <i>umbellatum</i>	30s.	<i>t</i> — <i>sperciliare</i>	"
<i>Cœlogyne</i> .		<i>t</i> — <i>venustum</i>	5-10s.
<i>t</i> — <i>barbata</i>	15s.	<i>t</i> — <i>villosum</i> (Illustr.)	15-25-50s.
<i>t</i> — <i>cristata</i>	10-50s.	<i>t</i> — <i>virens</i>	10s.
<i>t</i> — <i>elata</i>	15s.	Cyrtochilum , H. B. et K	
<i>t</i> — <i>flaccida</i>	10s.	<i>t</i> — <i>maculatum</i>	20s.
<i>t</i> — <i>flava</i>	25s.	Cyrtopodium	
<i>t</i> — <i>Massangeana</i>	500s.	<i>c</i> — <i>Andersoni</i>	10s.
<i>t</i> — <i>speciosa</i> (<i>Chelonanthera</i>).	20s.	<i>c</i> — <i>punctatum</i>	10s.
<i>t</i> — <i>testacea</i>	15s.	Dendrobium , Sw.	
<i>Colax</i> , Ldl.		<i>t</i> — <i>aemulum</i>	"
<i>t</i> — <i>jugosus</i> Ldl. (Illustr.)	20-75s.	<i>t</i> — <i>aggregatum</i>	15s.
<i>t</i> — de Puydti (Illustr.)	"	<i>t</i> — <i>majus</i>	"
<i>t</i> — <i>viridis</i>	30s.	<i>t</i> — <i>albo-sanguineum</i>	"
<i>Cymbidium</i> , Sw.		<i>t</i> — <i>aureum</i> (<i>heterocarpum</i>)	25-50s.
<i>t</i> — <i>albuciflorus</i>	20s.	<i>t</i> — <i>Bensoniae</i>	15-50s.
<i>t</i> — <i>eburneum</i>	15-200s.	<i>t</i> — <i>bigibbum</i> (Illustr.)	15s.
<i>t</i> — <i>giganteum</i>	25-75s.	<i>t</i> — <i>Boxallii</i>	"
<i>t</i> — <i>Lowi</i>	30-50s.	<i>t</i> — <i>Brymerianum</i>	"
<i>t</i> — <i>Mastersi</i>	20-50s.	<i>t</i> — <i>calceolare</i>	15s.
<i>Cypripedium</i> , Linn.		<i>t</i> — <i>Cambridgeanum</i>	15-35s.
<i>t</i> — <i>albo-purpureum</i>	"	<i>t</i> — <i>cariniferum</i>	30-50s.
<i>t</i> — <i>Argus</i>	20-50s.	<i>t</i> — <i>chrysanthum</i>	15s.
<i>t</i> — <i>Ashburtoniae</i>	38s.	<i>t</i> — <i>chrysotoxum</i> (III.)	15-30s.
<i>t</i> — <i>atro-purpureum</i>	10-25s.	<i>t</i> — <i>clavatum</i>	20-50s.
<i>t</i> — <i>barbatum</i>	5-10s.	<i>t</i> — <i>crassinode</i>	10-50s.
<i>t</i> — <i>grandiflorum</i>	15s.	<i>t</i> — <i>Barberianum</i>	25s.
<i>t</i> — <i>superbum</i>	10-25s.	<i>t</i> — <i>crepidatum</i>	10-30s.
<i>t</i> — <i>Boxalii</i> (Illustr.)	10-50s.	<i>t</i> — <i>crystallinum</i>	20-50s.
<i>t</i> — <i>Bullenianum</i>	25s.	<i>t</i> — <i>Dalhousianum</i> (III.)	20-50s.
* <i>t</i> — <i>caudatum</i> (<i>Selenipedium</i>) (<i>Pescat.</i>)	15-100s.	<i>t</i> — <i>Dayanum</i> (<i>anosmum</i>)	15-30s.
<i>t</i> — <i>caudatum roseum</i>	30-50s.	<i>t</i> — <i>densiflorum</i> (Illustr.)	15-25s.
<i>t</i> — <i>concolor</i> (Illustr.)	20s.	<i>t</i> — <i>Griffithi</i> (Illustr.)	20s.
<i>t</i> — <i>Dayanum</i>	50-150s.	<i>t</i> — <i>Devonianum</i> (Illustr.)	30s.
<i>t</i> — <i>Fairrieanum</i>	"	<i>t</i> — <i>eburneum</i>	15-20s.
<i>t</i> — <i>Hartwegi</i> (<i>Selenipedium</i>)	50s.	<i>t</i> — <i>Falconeri</i> (Illustr.)	15-100s.
<i>t</i> — <i>Haynaldianum</i>	30-100s.	<i>t</i> — <i>Farmeri</i> (<i>Pescat.</i>)	20-50s.
<i>t</i> — <i>Harrisianum</i>	25-50s.	<i>t</i> — <i>album</i>	"
<i>t</i> — <i>Hinksianum</i>	25-50s.	<i>t</i> — <i>fimbriatum</i>	10s.
<i>t</i> — <i>hirsutissimum</i>	25-75s.	<i>t</i> — <i>giganteum</i>	30-50s.
<i>t</i> — <i>Hookeri</i>	25s.	<i>t</i> — <i>oculatum</i> (Illustr.)	15s.
<i>t</i> — <i>insigne</i>	2-5s.	<i>t</i> — <i>formosum</i>	15-50s.
<i>t</i> — <i>Maulei</i>	5-10s.	<i>t</i> — <i>giganteum</i>	15-50s.
<i>t</i> — <i>javanicum</i>	15-30s.	<i>t</i> — <i>Freemannii</i>	15-25s.
<i>t</i> — <i>majus</i>	50s.	<i>t</i> — <i>Gibsonii</i>	15-25s.
<i>t</i> — <i>lævigatum</i>	50-100s.	<i>t</i> — <i>gracicaule</i>	15-25s.
<i>t</i> — <i>Lawrencianum</i>	60s.	<i>t</i> — <i>Griffithi</i>	25s.
* <i>t</i> — <i>longifolium</i> (<i>Selenip.</i>)	15-25s.	<i>t</i> — <i>Guiberti</i> (Illustr.)	"
<i>t</i> — <i>Lowi</i>	50-100s.		

<i>t</i> Dendrobium heterocarpum	15s.	Houilletia , A. Brogn
<i>t</i> — <i>Hilli</i>	25-50s.	<i>t</i> — <i>Brocklehurstiana</i> (Pescat.)
<i>t</i> — infundibulum (Illustr.)	20s.	15-30s.
<i>t</i> — <i>japonicum</i>	10s.	* <i>t</i> — <i>tigrina</i> (Illustr.) 30s.
<i>t</i> — <i>Kingianum</i>	25s.	Huntleya (Pescat.)
<i>t</i> — <i>macrophyllum</i> (<i>macran-</i> <i>thum</i>) (Pescat.)	15-50s.	<i>t</i> — <i>cerina</i> 25s.
<i>t</i> — — <i>giganteum</i>	40s.	Lælia , Ldl.
<i>t</i> — — <i>Dayanum</i>	25s.	<i>f</i> — <i>acuminata</i> 25s.
<i>t</i> — — <i>marmoratum</i>	10-25s.	<i>f</i> — <i>albida</i> 10-30s.
<i>t</i> — — <i>moniliforme</i>	10s.	<i>t</i> — <i>anceps</i> 10-25s.
<i>t</i> — — <i>monophyllum</i>	"	<i>t</i> — — <i>Barkeriana</i> 25s.
<i>t</i> — — <i>moschatum</i>	15s.	<i>t</i> — — <i>Dawsoni</i>
<i>t</i> — — <i>nobile</i>	10-25s.	<i>f</i> — <i>autumnalis</i> 25s.
<i>t</i> — — <i>Wallichii</i>	15-25s.	<i>f</i> — — <i>atrorubens</i> 50s.
<i>t</i> — — <i>nodatum</i>	15s.	<i>t</i> — — <i>Boothiana</i> 50s.
<i>t</i> — — <i>palpebrae</i>	20s.	<i>t</i> — — <i>cinnabarinia</i> 15-25s.
<i>t</i> — — <i>Parishi</i>	15s.	<i>t</i> — — <i>crispa</i> 10-25s.
<i>t</i> — — <i>Pierardi</i>	15s.	<i>t</i> — — <i>grandiflora</i> 25s.
<i>t</i> — — <i>latifolium</i>	"	<i>t</i> — — <i>Dayana</i> 25s.
<i>t</i> — — <i>Iutescens</i>	20s.	<i>t</i> — — <i>elegans</i> (Pescat.) 50-100s.
<i>t</i> — — <i>majus</i>	10-15s.	<i>t</i> — — <i>furfuracea</i> 15-25s.
<i>t</i> — — <i>primulinum</i>	15-25s.	<i>t</i> — — <i>majalis</i> 15s.
<i>t</i> — — <i>sanguinolentum purpureum</i>	25s.	<i>t</i> — — <i>Nelisi</i> (Illustr.) 200s.
<i>t</i> — — <i>secundum</i>	25s.	<i>t</i> — — <i>Perrini</i> 15-25s.
<i>t</i> — — <i>senile</i>	15-30s.	<i>t</i> — — <i>Pineli (præstans)</i> 25-50s.
<i>t</i> — — <i>speciosum</i>	15s.	<i>t</i> — — <i>purpurata</i> (Illustr.) 15-50s.
<i>t</i> — — <i>suavissimum</i>	20-50s.	<i>t</i> — — <i>Brysiana</i> 15-50s.
<i>t</i> — — <i>sulcatum</i>	30s.	<i>t</i> — — <i>Schilleriana splendens</i> 150s.
<i>t</i> — — <i>teretefolium</i>	"	<i>f</i> — <i>superbiens</i> 20s.
<i>t</i> — — <i>tetragonum</i>	30s.	<i>t</i> — — <i>xanthina</i> 15-30s.
<i>t</i> — — <i>thyrsiflorum</i> (III.)	15-50s.	Leptotes , Ldl.
<i>t</i> — — <i>transparens</i>	15s.	<i>t</i> — — <i>bicolor</i> 10-25s.
<i>t</i> — — <i>Wardianum</i> (III.)	15-100s.	Limatodes , Bl.
Dendrochilum , Bl.		<i>t</i> — — <i>rosea</i> 10-25s.
<i>t</i> — — <i>glumaceum</i>	20s.	Miltonia , Ldl.
<i>t</i> — — <i>latifolium</i>	25s.	<i>t</i> — — <i>candida</i> 20s.
Epidendrum , Linn.		<i>t</i> — — var. <i>Yenischiana</i> 50s.
<i>t</i> — — <i>ambiguum</i> (Illustr.)	15s.	<i>t</i> — — <i>cereola</i> 30-50s.
<i>t</i> — — <i>aurantiacum</i>	15s.	<i>t</i> — — <i>Clowesi</i> 10-30s.
<i>c</i> — <i>bicornutum</i>	20s.	<i>t</i> — — <i>Moreliana</i> (Illustr.) 25-50s.
<i>t</i> — — <i>Brasavolæ</i>	30s.	<i>t</i> — — <i>Regnelli</i> (Illustr.) 30-50s.
<i>t</i> — — <i>ciliare</i>	10s.	<i>t</i> — — <i>purpurea</i> 75s.
<i>t</i> — — <i>cochleatum macrobulbon</i>	10s.	<i>t</i> — — <i>spectabilis</i> 10-25s.
<i>t</i> — — <i>fragrans</i>	10s.	* <i>t</i> — — <i>virginialis</i> (Illustr.) 20s.
<i>t</i> — — <i>glumaceum</i>	25s.	<i>t</i> — — <i>Warscewiczi</i> (III.) 20-50s.
<i>t</i> — — <i>macrochilum</i>	10-25s.	Oncidium , Sw.
* <i>t</i> — — <i>pentotodes</i> , Rchb. f.	25s.	— <i>ampliatum majus</i> 20s.
<i>t</i> — — <i>Stamfordianum</i>	20-50s.	<i>t</i> — — <i>Batemani</i> 15-25s.
Esmeralda (<i>Vanda</i>), Rchb. f.		<i>f</i> — <i>bifolium</i> 25s.
<i>t</i> — — <i>Cathcarthi</i> (Illustr.)	50-150s.	<i>t</i> — — <i>Carthaginense</i> 10s.
Galeandra Devoniana (III.)	25s.	<i>t</i> — — <i>Cavendishianum</i> 15s.
Gongora , R. et Pav.		<i>t</i> — — <i>celbolleta (juncifolium)</i> 10-20s.
<i>t</i> — — <i>armeniaca</i>	15s.	* <i>t</i> — — <i>exasperatum</i> 20-50s.
<i>t</i> — — <i>leucochila</i>	25s.	<i>t</i> — — <i>flexuosum</i> 10s.
* <i>t</i> — — <i>odoratissima</i>	15s.	<i>t</i> — — <i>fuscatum (Miltonia Wars-</i>
* <i>t</i> — — <i>portentosa</i> (Illustr.)	25s.	<i>zewiczi</i>) (Illustr.) 15-50s.
* <i>t</i> — — <i>scaphephorus</i>	15s.	<i>t</i> — — <i>gyrobolbon</i> 40s.
Grammatophyllum , Bl.		* <i>t</i> — — <i>micranthum</i> 25s.
— <i>Ellisi</i>	35s.	<i>t</i> — — <i>janeirensis</i> (Illustr.) 10s.
		<i>t</i> — — <i>juncifolium</i> 10s.
		* <i>t</i> — — <i>Kramerianum</i> 20s.

<i>c</i> <i>Oncidium</i> <i>Lanceanum</i>	20-50s.	<i>Stanhopea</i> <i>Devonian</i>	25s.
<i>c</i> — <i>luridum</i>	15s.	— <i>Dowiana</i>	"
<i>c</i> — <i>guttatum</i>	25s.	— <i>eburnea</i>	"
<i>t</i> — <i>Marshallianum</i>	"	— <i>grandiflora</i>	"
* <i>t</i> — <i>macranthum</i>	"	<i>t</i> — <i>Haseloffi</i>	25s.
<i>t</i> — <i>Papilio</i>	10-25s.	— <i>insignis</i>	"
<i>t</i> — <i>major</i> (<i>Illustr.</i>)	20s.	— <i>Martiana</i>	"
<i>t</i> — <i>pubes</i>	15s.	<i>t</i> — <i>oculata</i>	50s.
<i>t</i> — <i>pulvinatum</i>	10s.	<i>t</i> — <i>tigrina</i>	25s.
<i>t</i> — <i>ramosum</i>	20s.	<i>Trichocentrum</i> , Poepp. et Endl.	
<i>t</i> — <i>reflexum</i>	25s.	<i>t</i> — <i>tigrinum</i> (<i>III.</i>)	75s.
— <i>sarcodes</i> (<i>Illustr.</i>)	20s.	<i>c</i> — <i>albo-coccineum</i>	"
<i>t</i> — <i>sphacelatum</i>	10-25s.	<i>t</i> — <i>Pfavi</i>	50s.
<i>t</i> — <i>stelligerum</i>	20s.	<i>c</i> — <i>roseum</i>	75s.
<i>t</i> — <i>Wentworthianum</i>	10-25s.	<i>Trichopilia</i> , Ldl.	
<i>Paphinia</i> <i>cristata</i> Ldl.	"	<i>t</i> — <i>coccinea</i> (<i>marginata</i>)	15-30s.
<i>Peristeria</i> , Hook.		<i>t</i> — <i>fragans</i> (<i>Pilumna</i>)	25s.
<i>t</i> — <i>clata</i>	25-30s.	<i>t</i> — <i>suavis</i>	15-25s.
<i>Pescatoria</i> , Rehb. f.		<i>t</i> — <i>tortilis</i>	10-25s.
<i>t</i> — <i>Bruchmüller</i>	50s.	<i>Uropedium</i> , Ldl.	
<i>t</i> — <i>cerina</i>	40-60s.	* <i>t</i> — <i>Lindeni</i> (<i>Pescat.</i>)	25-50s.
<i>t</i> — <i>Lehmanni</i> (<i>Illustr.</i>)	"	<i>Vanda</i> , R. Br.	
<i>t</i> — <i>Roezli</i>	30-50s.	<i>c</i> — <i>Batemani</i>	25-500s.
<i>Phajus</i> , Lour.		<i>c</i> — <i>Bensoni</i>	50s.
<i>t</i> — <i>grandiflorus</i>	5s.	<i>t</i> — <i>Carthcarti</i>	50-1500s.
<i>t</i> — <i>maculatus</i>	10s.	<i>t</i> — <i>cœrulea</i> (<i>Pescat.</i>)	50s.
<i>t</i> — <i>Wallichii</i>	15s.	<i>c</i> — <i>Denisoniana</i> (<i>Illustr.</i>)	50s.
<i>Phalaenopsis</i> , Bl.		<i>c</i> — <i>gigantea</i>	20-50s.
<i>c</i> — <i>amabilis</i>	25-150s.	<i>c</i> — <i>Hookeri</i> (<i>Illustr.</i>)	20s.
<i>c</i> — <i>Esmeralda</i>	25s.	<i>c</i> — <i>insignis</i>	75-200s.
<i>c</i> — <i>equestris</i> (<i>rosea</i>)	35-75s.	— <i>lamellata</i>	100s.
<i>c</i> — <i>grandiflora</i>	20-100s.	<i>c</i> — <i>Lowi</i> (<i>Illustr.</i>)	150-500s.
<i>c</i> — <i>Luddemanniana</i>	40-100s.	<i>c</i> — <i>Roxburghii</i>	30s.
* <i>c</i> — <i>Schilleriana</i> (<i>Illustr.</i>)	20-50s.	<i>c</i> — <i>cœrulea</i>	50s.
<i>c</i> <i>Renanthera</i> <i>coccinea</i>	25s.	<i>c</i> — <i>suavis</i> (<i>Pescat.</i>)	50-200s.
<i>c</i> — <i>Lowi</i> (<i>Vanda</i>)	150s.	<i>c</i> — <i>teres</i>	10-25s.
<i>Saccolabium</i> , Bl.		<i>c</i> — <i>tricolor</i> (<i>Pescat.</i>)	25-100s.
<i>c</i> — <i>ampullaceum</i>	15-50s.	<i>e</i> — <i>aurea</i>	50-150s.
<i>c</i> — <i>Blumei</i>	15-75s.	<i>c</i> — <i>— cinnamomea</i>	30-150s.
<i>c</i> — <i>— majus</i> (<i>Illustr.</i>)	10-125s.	<i>c</i> — <i>— Leopoldi</i>	75s.
<i>c</i> — <i>curvifolium</i>	20-50s.	<i>c</i> — <i>— Loddigesii</i>	75s.
<i>t</i> — <i>densiflorum</i>	20s.	<i>c</i> — <i>— formosa</i> (<i>V. suavis</i> <i>Rollisson</i>)	50-150s.
<i>c</i> — <i>giganteum</i>	20-200s.	<i>c</i> — <i>tricolor</i> <i>purpureo-mar-</i>	
<i>c</i> — <i>— longifolium</i>	75-200s.	<i>ginata</i>	75s.
<i>c</i> — <i>guttatum</i>	25-200s.	<i>c</i> — <i>— Schilleriana</i>	75-150s.
<i>c</i> — <i>Harrisonianum</i>	75s.		
— <i>illustre</i>	50-100s.	<i>*c</i> — <i>cyanæa</i>	25s.
<i>c</i> — <i>premorsum</i>	50-90s.	<i>Warscewiczella</i> , Rohb.	
<i>c</i> — <i>violaceum</i>	25-100s.	<i>t</i> — <i>discolor</i>	20-50s.
<i>Sarcopodium</i> , Ldl.		<i>Zygotepetalum</i> , Hook.	
<i>t</i> — <i>Lobbi</i>	15-30s.	<i>t</i> — <i>crinitum</i>	25s.
<i>Schomburgkia</i> , Ldl.		* <i>t</i> — <i>Gautieri</i>	30-100s.
* <i>t</i> — <i>gloriosa</i>	50s.	* <i>c</i> — <i>graminifolium</i>	50s.
* <i>t</i> — <i>rosea</i>	25s.	<i>t</i> — <i>Mackayi</i>	10-15s.
<i>Scuticaria</i> , Ldl.			
<i>c</i> — <i>Steeli</i>	50s.		
<i>Stanhopea</i> , Hook.			
<i>t</i> — <i>Bucephalus</i>	10s.		
<i>t</i> — <i>cinnamomiodora</i>	15s.		
<i>t</i> — <i>cirrhata</i>	15s.		

The special list of large and choice specimens will be sent on application.

COLLECTIONS OF ORCHIDS.

1st Collection

12 Coolhouse varieties for 200s.

Ada aurantiaca.
Odontoglossum Alexandræ.
 " *Pescatorei.*
 " *triumphans.*
 " *cirrhosum.*
Masdevallia amabilis.
 " *Lindeni.*
Lycaste Skinneri.
Oncidium incurvum.
 " *tigrinum.*
 " *crispum.*
Lælia affine furfuracea.

3rd Collection

12 Coolhouse varieties for 100s.

Lycaste Skinneri.
 " *Harrissoniana.*
Lælia affine furfuracea.
Oncidium crispum.
 " *incurvum.*
 " *Barkeri.*
Odontoglossum Alexandræ.
 " *triumphans.*
Epidendrum vitellinum majus.
Zygopetalum cœrulescens.
Masdevallia civilis.
Dendrobium nobile

5th Collection

25 varieties for 150s.

Aerides odoratum.
Odontoglossum bictoniense.
 " *Rossi majus.*
 " *maculatum.*
Oncidium sphacelatum.
 " *crispum.*
 " *roseum.*
 " *papilio.*
Cypripedium barbatum grandiflor.
 " *Roezli.*

2nd Collection

12 Warmhouse varieties for 200s.

Cypripedium Stonei.
 " *Hookeri.*
 " *niveum.*
Dendrobium formosum.
 " *densiflorum.*
Miltonia Clowesi.
Aerides odoratum.
Vanda tricolor.
 " *Roxburghii.*
Phalænopsis amabilis.
 " *Schilleriana.*
Cattleya Harrissoniana.

4th Collection

12 Warmhouse varieties for 150s.

Aerides virens Dayi.
 " *crassifolium.*
 " *odoratum.*
Vanda tricolor.
Oncidium papilio.
Phalænopsis amabilis.
 " *Schilleriana.*
Saccolabium Blumei majus.
Schomburgkia rosea.
Zygopetalum Makavi.
 " *graminifolium.*
Lælia purpurata.

6th Collection

10 Selected varieties for 50s.

Masdevallia tovarensis.
Oncidium tigrinum.
Cypripedium barbatum.
 " *insigne.*
Epidendrum macrochillum.
Schomburgkia rosea.
Cattleya Mossiae.
Odontoglossum maculatum.
Eria convallarioides.
Oncidium crispum.

Dendrobium nobile.
 " fimbriatum.
 Chysis aurea.
 Cattleya Harrissoniana.
 " Mossiae.
 Lycaste Deppei.
 " Harrissoniana.
 Oncidium pubes.
 " ampliatum.
 " Barkeri.
 Masdevallia civilis.
 " polysticta.
 Stanhopea tigrina.
 Lælia pedoncularis.
 Schomburgkia gloriosa.

7th Collection**10 Coolhouse varieties for 50s.**

N. B. These are winter-flowering and especially suitable for bouquet-making.

Masdevallia tovarensis.
 Odontoglossum Alexandræ.
 " Pescatorei.
 " cordatum.
 Lycaste Skinneri.
 Sophronitis grandiflora.
 Cypripedium insigne.
 " barbatum.
 Cœlogyne cristata.
 Dendrobium nobile.

CONSERVATORIES. — The Establishment possesses a very considerable selection of Palms, Tree Ferns and other plants in vigorous specimens for the ornamentation of *Conservatories* and large stoves, etc., of which a list will be communicated on application being made, stating species, sizes and prices. *The firm undertakes the furnishing of Conservatories* and has persons especially devoted to this branch of work.

TREE FERNS.

Very complete collection, varying from the largest species to the smallest, such as the *Todea Wilkesiana*.

Prices of specimens on application.

<i>Alsophila australis</i>	15-30-100s.
— <i>Beyrichiana</i>	— —
— <i>Colensoi</i>	— —
— <i>contaminans (glauca)</i> (Illustr.)	500s
— <i>excelsa</i>	100-300s.
— <i>Leichardtiana</i>	— —
— sp. <i>Sanctæ Catherine</i>	— —
— <i>Paraguayensis</i>	150-300s.
<i>Balantium antarcticum</i>	30-75-150s.
<i>Blechnum brasiliense</i>	25-75-150s.
— <i>corcovadense</i>	— —
<i>Cibotium princeps</i>	5-1500s.
— <i>australis (hybrid.)</i>	1800s.
— <i>Schiedei</i>	3-1200s.
— <i>spectabile</i>	3-1200s.
<i>Cyathea Beirichiana</i>	3-1000s.
— <i>Blanchei</i>	— —
— <i>Burkei</i>	200-300s.
— <i>Cunninghamii</i>	100-250s.
— <i>dealbata</i>	100-350ss
— <i>Dregei</i>	200-350s.
— <i>funebris</i>	— —

Cyathea medullaris	25-75-150-500s.
— paraguayensis	100-150s.
— Smithii	150-300s.
Dicksomia antarctica	25-50-75-150s.
— <i>Deplanchei</i>	100-300s.
— lanata	5-800s.
— Smithii	3-1000s.
— squarrosa	30-75-150s.
— Youngii	—
Didymochlaena truncata	25-100s.
Hemitelia guyanensis	100-300s.
— horrida	125-300s.
— Smithii	—
Lomaria Boryana	50-100s.
— cycadæfolia	20-50s.
— <i>gigantea</i> (<i>Illustr.</i>)	30-150s.
— neo-caledonica (<i>Illustr.</i>)	—
Todæa barbara	3-800s.
— Fraserii	15s.
— intermedia	15s.
							Very large.	200s.
— Vroomi	15s.
— Wilkesiana (<i>Illustr.</i>)	100s.

Angiopteris australis	20-250s.
— erecta	5-15-800s.
Marattia attenuata (<i>Illustr.</i>)	200-800s.
— furluracea	50-600s.
— Laucheana	—
— macrophylla	25-300s.
— Verschaffelti	25-300s.

C R O T O N.

Croton albicans	.	20s.		Croton crysophyllum	.	.	.	5-15s.
— Andreanum (<i>Illustr.</i>)	.	3-5-10s.		— Cooperii	.	.	.	5-10s.
— angustifolium	.	2-5s.		— cornutum	.	.	.	2-15s.
— Anatomensis	.	5-10s.		— Crown Prince	.	.	.	25s.
— appendiculatum	.	5-10s.		— discolor	.	.	.	5s.
— Aucubæfolium	.	2s.		— d'Israëli	.	.	.	5-10s.
— Baronne James de Roth-				— Dormanianum	.	.	.	25s.
schild (<i>Illustr.</i>)	.	5-50s.		— Donneai	.	.	.	10s.
— Bergmannii (<i>Illustr.</i>)	.	10-50s.		— Earl of Derby	.	.	.	5-10s.
— Bismarcki	.	5-10s.		— elegantissimum	.	.	.	5-10s.
— Burtoni	.	10s.		— Evansianum	.	.	.	10-25s.
— campophyllum	.	5-10s.		— faleatum	.	.	.	5-10s.
— Carrieri	.	10-50s.		— fasciatum	.	.	.	5-15s.
— Challenger	.	10s.		— Fenzi	.	.	.	35s.
— Chelsoni	.	15-25s.		— fuscatum	.	.	.	2s.
				— Goedenoughti	.	.	.	5-10s.

Croton Goldiei	15s.	Croton mutabilis	15s.
— grande	3s.	— nobilis	10s.
— guineensis	80s.	— nigrum	5s.
— hastiferum (<i>Illustr.</i>)	5-10s.	— ovalifolium	5-10s.
— Hendersoni.	5-10s.	— paradoxum	15s.
— Henryanum	20s.	— pictum (<i>Illustr.</i>)	2-50s.
— Hookeri	5-10s,	— picturatum	15s.
— Hillianum	3s.	— Prince of Wales	10s.
— hybridum	5-10s.	— Queen Victoria	5-10s.
— imperator	10s.	— reginae	15s.
— intermedium	5-10s.	— Rex	5-10s.
— irregulare	2-5s.	— roseo-pictum (<i>Illustr.</i>)	10s.
— Jamesi	8-15s.	— Royal Prince	5-10s.
— Johannis	3-10s.	— Sinitzini	10s.
— lacteum	3-10s.	— Spinckii	15s.
— longifolium	2s.	— spirale	5-10s.
— limbatum	6s.	— Stræhmeri	25s.
— Lord Cairns	10-25s.	— sunset	10s.
— lyratum (<i>Illustr.</i>)	5-15s.	— torquatum	10s.
— Macafeanum	10-25s.	— tortile	15s.
— Macfarlanei	20s.	— trilobum	5-10s.
— maculatum Katoni	5s.	— Truffateanum	20-150s.
— magnificum (<i>Illustr.</i>)	15s.	— undulatum	3-10s.
— majesticum	5-10s.	— variegatum	2s.
— Massangeanum (<i>Illustr.</i>)	10-50s.	— Veitchii	2-5s.
— Mc Arthurii	10-20s.	— Vervaeti (<i>Illustr.</i>)	10-25s.
— microphyllum	5-10s.	— volutum	5-10s.
— Mooreanum	5-10s.	— Weismanni	3-5s.
— multicolor	3-5s.	— Williamsii	15s.

Collection of 12 Crotons, our selection : 18, 50, 100 shillings.

 The price of Show plants will be forwarded on application being made.

L'ILLUSTRATION HORTICOLE

L'ILLUSTRATION HORTICOLE, monthly review of gardens and flowers.

For conditions of subscription see page 2 of the cover.

DRACÆNA.

 The best varieties are printed in larger type.

Dracæna albicans	3-5s.	Dracæna Leopoldi	10-30s.
— albo marginata	6-30s.	— Levangeri	5-10s.
— amabilis (Illustr.)	3-10s.	— Liervali	10s.
— Anerleyensis	5-30s.	— limbata	2-25s.
— angustifolia	10s.	— Lindenii (Illustr.)	50s.
— atro-purpurea	30s.	— longifolia alba	50s.
— angusta	5-10s.	— lutescens	10s.
— australis	2-10s.	— Macleayi	3-5s.
— Balmoreana	5-10s.	— macrophylla	30s.
— Banksii	3-10s.	— M ^e Lecocq Dumenil	20s.
— Baptisti (Illustr.)	3-15s.	— majestica	20-50s.
— Baroni	5-25s.	— magnifica	3-10s.
— Bausei	10-25s.	— Massoni	10-30s.
— Bergmanni	20s.	— mauritiana	5s.
— brasiliensis	2-10s.	— metallica	10-25s.
— cannefolia	2-10s.	— Mr Charles Heine	30s.
— Cantrelli	5-20s.	— Moorei	3-10s.
— Casanovæ (Illustr.)	10-15s.	— Mosevi	30s.
— Chantrieri	50s.	— Mrs C. J. Freake	10-25s.
— Chelsoni	5-10s.	— Neo-Caledonica (Illustr.)	10-25s.
— Cooperi	2s.	— nigricans	2s.
— compacta	5-10s.	— nigro-rubra	2s.
— concinna	5-10s.	— Nitzscherii	5-25s.
— congesta	1s.	— oculata	5s.
— Dennisoni	5-10s.	— ornata	8-20s.
— Draco	2s.	— pendula	2-5s.
— Duffi	10s.	— picta	10s.
— Elisabethæ	5-10s.	— porphyrophylla	3-10s.
— erecta alba (Illustr.)	50s.	— princeps	5-10s.
— ferrea	3-5s.	— purpurea	30s.
— Fraseri	10s.	— excelsa	3-10s.
— fructuosa	2-10s.	— pulchella	5-10s.
— Gladstoni	5-20s.	— pulcherrima	5-10s.
— gloria (Illustr.)	3-10s.	— punctata	3s.
— Goldieana (Illustr.)	10-30s.	— Reali (Illustr.)	10-5s.
— grandis	5-10s.	— Rebecca	10-25s.
— graezielii (Illustr.)	2s.	— Robinsoniana (Illustr.)	10-25s.
— Guilfoylei (Illustr.)	2-5s.	— recurva	10s.
— Haageana	3s.	— recurvata	30s.
— Hibberti	1s.	— regalis	10s.
— hybrida	5-10s.	— Reginæ	3s.
— illustris	15-25s.	— regis (Illustr.)	75s.
— imperialis	2-15s.	— robusta	2-15s.
— imperator	10-25s.	— rubella	10s.
— indivisa	2-5s.	— Rumphi	3-5s.
— lentiginosa	10s.	— salmonea	10-30s.
— lineata	2-5s.	— siamensis	4-10s.
— ignea	20s.	— splendens	3s.
— jaspidea (Illustr.)	5-10s.	— superba	20s.
— Knerkiana	2-5s.	— Sydneyi	10-25s.
— latifolia (grandis)	10-25s.	— Taylori	25s.

Dracæna Tellingii	5-25s.	Dracæna vigidiifolia	2s.
— <i>terminalis</i>	3-10s.	— <i>Warocquei</i> (<i>Illustr.</i>)	10-25s.
— — <i>alba</i>	5-30s.	— <i>Weismanni</i>	5s.
— Thompsoni	10-25s.	— <i>Wilsoni</i>	10-30s.
— <i>umbraculifera</i>	2-5s.	— Youngi	2-10s.
— <i>Verloti</i>	10-25s		

All the varieties named above are available in specimens and half specimens for exhibitions.

12 Fine varieties our selection 25, 50, 100, and 200 shillings.

BROMELIADS.

We can supply fine collections of 12 varieties our selection for 30, 50, 150s. according to force.

Aechmea cærulescens	3-10s.	Billbergia pyramidalis	5s.
— <i>cœlestis</i>	2s.	— <i>Quesneliana</i>	10-25s.
— <i>distiacantha</i>	5-10s.	— <i>Reginæ Ameliæ</i>	15s.
— <i>fulgens</i>	3-5s.	— <i>rhodocyanea</i>	5s.
— — <i>discolor</i>	3-5s.	— — <i>discolor</i>	10s.
— <i>Glaziovii</i>	1s.	— <i>rosea</i>	10s.
— <i>Luddemaniana</i>	10s.	— <i>sanguinea</i>	3s.
— <i>Maria Reginæ</i>	5-25s.	— <i>splendida</i>	2-5s.
— <i>miniata</i>	5-10s.	— <i>superba</i>	10s.
— <i>spectabilis</i>	3-10s.	— <i>thyrsoidæa</i>	3s.
— <i>Veitchi (Chevalieri)</i>	15-50s.	— <i>Wiotiana</i>	3s.
— <i>Weilbachi</i>	3-10s.	— <i>Zebrina</i>	2-5s.
Ananassa Bracamorensis	50s.		
— <i>Mordilona</i>	25-50s.	Bromelia Binoti	5s.
— <i>pinangensis</i> fol. varg.	40s.	— <i>bracteata</i>	25s.
— <i>Porteana</i>	15s.	— <i>Karatas</i>	4s.
— <i>sativa</i> fol. var.	20-40s.	— <i>Pingoin</i>	4-10s.
12 Ananas variés.	75-120-300s.	— <i>scæptrum</i>	10s.
Anoplophyton strictum	15s.	— <i>undulatum</i>	3-10s.
Billbergia amœna	3s.	Canistrum aurantiacum	5s.
— <i>clorosticta</i>	10s.	— <i>splendens</i>	3-10s.
— <i>discolor</i>	4s.	Caraguata cardinalis (<i>Illustr.</i>)	50-100s.
— <i>excellens</i>	10s.	— <i>discolor</i>	5-10s.
— <i>farinosa</i>	5s.	— <i>Van Volxemi</i> (<i>Illustr.</i>)	20-100s.
— <i>fasciata</i>	5s.	Chevalliera Veitchi	15-50s.
— <i>gigantea</i>	10-25s.	Cryptanthus acaulis	3-5s.
— <i>Joinvillei</i>	2s.	— <i>brevittatus</i>	3-5s.
— <i>Leopoldi</i>	3-5s.	— <i>clavatus</i>	3-5s.
— <i>Liboniana</i>	3-5s.	— <i>fusces</i>	3s.
— <i>longifolia</i>	2s.	— <i>longifolius</i>	5s.
— <i>melanacantha</i>	4s.	— <i>zonatus</i>	3-5s.
— <i>Moreliani</i>	5s.	Disteganthus basilateralis	3-10s.

Disteganthus Moensi	250s.	Pitcairnea imbricata	4s.
— scarlatinus	20s.	— irridifolia	5s.
Echinostachys Pineliana	10s.	— polyanthoides	4s.
Encholirion corallinum	3-10s.	— rivalis	15s.
— Jonghi	3-10s.	— speciosissima	4s.
— roseum	3-10s.	— staminea	4s.
— sanguinolentum	25s.	— sulphurea	4s.
— Saundersi	3-10s.	— tabulaeformis	3-5s.
Greigia sphacelata	4s.	— undulatofolia	10s.
Guzmania erthroplepis	19s.	Portea kermesina	10-25s.
— fragrans	4-10s.	Pourretia Achupalla	10s.
— grandis	10s.	— brevifolia	3-15s.
— Melinonis (new)	15s.	— mexicana	4s.
— picta	6s.	— yuccoides	15s.
— Reginæ	15-60s.	Puya gigantea	10s.
— tricolor	3-6s.	— heterophylla	5s.
Hechtia argentea	3-10s.	— recurvata	5s.
— Ghiesbreghtii	3-10s.	Quesnelia rufa	10-25s.
— glomerata	15s.	Ronnbergia Morreniana (III.)	15-50s.
— pitcairniaefolia	3-10s.	Ruyschia Ellemetti	10s.
Helicodia Baraquimiana	5s.	Tillandsia bivittata	5s.
Hohenbergia erythrophachys	10s.	— complanata	10-25s.
— strobilacea	2-10s.	— cyanea	5s.
Hoplophyton calyculatum	10s.	— dianthoides	1' s.
— giganteum	10s.	— farinosa	2s.
— Lindeni	10s.	— fasciata	20s.
Macrocordium tinctorium	10s.	— glaucophylla	8s.
Massangea musaica	30-75s.	— guttata	15-30s.
Nidularium amazonicum	5-25s.	— heterophylla	10s.
— agavefolium	5s.	— juncifolia	30s.
— Bahiense	10s.	— Legrellei	5-10s.
— coriaceum	10-20s.	— Lindeni	25-100s.
— fulgens	6s.	— — vera (Illustr.)	50-100s.
— atro-purpureum	5-10s.	— — musaica (Illustr.)	30-75s.
— discolor	3s.	— pruinata	25-50s.
— Innocenti	3-5s.	— Rollisoni	15s.
— maculatum	10s.	— splendens	2-5s.
— marmoratum	10s.	— — major	15s.
— Meyendorffii	3-10s.	— stricta	25s.
— Morrenianum	25s.	— tessellata (Illustr.)	30-50s.
— pictum	5-10s.	— xiphioides	10s.
— princeps	10s.	— Zahni	2's.
— purpureum	2-5s.	— zebrina	2-5s.
— Scheremetieffii	3s.	Tussaccia nitida	10s.
— spectabile	10s.	Vriesea brachystachys	5s.
— triste	2s.	— glaucophylla	10s.
Ortglesia palliolata	5-10s.	— Glaziovana	5-15s.
— tillandsioides	5-10s.	— Hillegeeriana	25s.
Pepinia aphelandraflora	2-5s.	— Malzini	5-10-25s.
— iridifolia	3-5s.	— princeps	25s.
— punicea	2s.	— psittacina	5-10s.
Pholidophyllum zonatum	3s.	— sanguinolenta	50s.
Pitcairnea corallina	5-10s.	— splendens	5s.

Specimens are available. Collection of 12 Bromeliads for exhibiting,
100, 250 shillings and upwards.

ECONOMIC AND MEDICINAL PLANTS.

Economic and Medicinal Plants form one of the specialities of the establishment. We forward them to all parts of the world employing such a system of packing as ensures their safe arrival.

We would call the attention of all *Botanic Gardens* to the fact that we are always willing to exchange plants whether new or rare.

<i>Abrus precatorius</i>	<i>Crabs Eyes</i>	15s.
<i>Adenanthera pavonina</i> .	<i>Barricari Seeds</i>	5s.
<i>Albizia Lebbecki</i>		10-20s.
<i>Alibertia (Gardenia) edulis.</i>		25s.
<i>Amomum Cardamomum</i>		1s.
<i>Amyris balsamifera</i>		25s.
<i>Anda Gomezi</i>		15s.
<i>Antiaris toxicaria</i> . <i>Upas tree</i>		25s.
<i>Arenga saccharifera</i> . <i>Suyar Palm</i>		25s.
<i>Barringtonia speciosa</i>		25s.
<i>Bassia latifolia</i> . <i>Mahva Tree</i>		15s.
<i>Bauhinia multinervia</i>		25s.
<i>Bixa Orellana</i> . <i>Dye Plant</i>		5s.
<i>Bombax Ceiba</i> . <i>Silk Cotton Tree</i>		10s.
<i>Brachychiton populneum</i> . <i>Bottle Tree</i>		5s.
<i>Cæsalpinia echinata</i>		5s.
<i>Calophyllum Calaba</i> . <i>Calaba Oil</i>		10s.
— <i>inophyllum</i>		10s.
<i>Canella alba</i> . <i>Canella Bark</i>		15s.
<i>Carapa guyanensis</i>		5s.
<i>Carludovica palmata</i> . <i>Panama straw</i>		5-10s.
<i>Cassia fistula</i>		10s.
— <i>glutinosa</i>		10s.
— <i>moschata</i>		10s.
<i>Castilloa elastica</i> . <i>Panama India rubber</i>		25s.
<i>Cecropia palmata</i>		2s.
<i>Cedrela brasiliensis</i> . <i>Cedar wood</i>		15s.
— <i>odorata</i>		5-10s.
<i>Cephaëlis Ipecacuanha</i> . <i>Ipecacuanha</i>		10s.
— <i>mucosa</i>		10s.
<i>Cerbera tanguin</i> . <i>The tanguin, poison of Madagascar</i>		15s.
<i>Ceroxylon andicola</i> . <i>Wax Palm</i>		15s.
<i>Chavica Betel</i> . <i>Betel Nut</i>		10s.
<i>Chloroxylon Swietenia</i> . <i>Satin wood</i>		10s.
<i>Cinchona officinalis</i> . <i>Crown bark</i>		10s.
— <i>pubescens</i>		15-25s.
— <i>succi-rubra</i> . <i>Red bark</i>		10s.

Cinnamomum aromaticum.	5s.
— <i>dulce</i>	2s.
— <i>sericeum</i>	5s.
— <i>zeylanicum. Ceylon cinnamon.</i>	5s.
Cissampelos Pareira <i>Pareira brava</i>	10s.
Citrosma Lindeni	10s.
— <i>Thea</i>	15s.
Coffea arabica. Coffee	15s.
— — <i>augustifolia</i>	2-10s.
— — <i>bengalensis</i>	10s.
— — <i>Moka. Moka Coffee.</i>	5-10s.
— — <i>myrtiflora</i>	10s.
— — <i>liberiana. Liberian Coffee</i>	15-25s.
Colocasia esculenta	3s.
Condaminea longifolia. Febrifuge	26s.
— — <i>macrophylla. Febrifuge</i>	25s.
Crescentia Cujete. Calabash tree.	5s.
— — <i>regalis</i>	5s.
Croton Cascarilla. Croton Oil Plant	5s.
Curcas purgans (Jatropha)	5s.
Dalbergia Sissoo	10s.
Dillenia speciosa	10s.
Diospyros Ebenum. Ceylon ebony	15s.
Elaeodendron australe. Olive wood	15s.
Erythoxylon Coca (Illustr.). Coca	10s.
— — <i>macrophyllum</i>	10s.
Euterpe edulis. Cabbage Palm.	5s.
— — <i>oleracea</i>	10s.
Galipia odoratissima. Febrifuge	5s.
Geoffroya vermiculata (Andira anthelmintica)	15s.
Gossypium arboreum. Cotton Tree	5s.
Guayacum arboreum.	15s.
— — <i>officinale Lignum Vitæ</i>	15s.
Gynerium saccharoides	5s.
Hæmatoxylon campechianum. Logwood	5s.
Herietiera littoralis	10s.
Hermandia peltata	10s.
Hura crepitans. Sandbox tree	10s.
Hymenæa Courbaril. Locust tree	5-10s.
Ilex Congoin. Maté de St-Paul	5s.
— — <i>paraguayensis. Paraguay-Tea</i>	5s.
Inga laurina	10s.
Janipha Manihot. Mandioca	5s.
Jatropha Cureas	5s.
Lagetta funifera	15s.
Laurus Benzoin.	5s.
— — <i>Camphora. Camphor</i>	5-10s.
— — <i>Cannella</i>	5s.
— — <i>cinnamomum</i>	5s.

<i>Maba seychellarum</i>	10s.
<i>Macadamia ternifolia</i>	<i>Queensland Nut</i>	25s.
<i>Machærium firmum</i>	<i>Rosewood</i>	5-10s.
<i>Macropiper excelsum</i>	5s.
<i>Manihot utilissima</i>	<i>Bitter Cassava</i>	5s.
<i>Maranta arundinacea</i>	<i>Arrow root</i>	5s.
<i>Mimosa vaga</i>	10s.
<i>Mimusops Balota</i>	<i>Gutta-Percha</i>	10s.
<i>Monodora grandiflora</i>	25s.
<i>Murraya exotica</i>	5s.
<i>Musa textillis</i>	25s.
<i>Mussænda frondosa</i>	10s.
<i>Myrcia pimentoïdes</i>	10s.
<i>Myristica tubiflora</i>	10s.
—	<i>moschata. Nutmeg.</i>	5s.
<i>Myrospermum peruiferum</i>	10s.
<i>Myroxylon Pereira</i>	5s.
<i>Ochromia borbonica</i>	10s.
<i>Oldfieldia africana</i>	10s.
<i>Pandanus utilis. Vauquois</i>	3-10s.
<i>Phytelephas macrocarpa. Vegetable Ivory</i>	15s.
<i>Pimenta vulgaris. Red pepper.</i>	5s.
—	<i>acris. Wild Clove</i>	5s.
—	<i>officinalis. Allspice.</i>	5s.
<i>Piper Betel. Betle pepper leaf.</i>	5s.
—	<i>Futocadsura. Japanese pepper</i>	3s.
—	<i>longum.</i>	5s.
—	<i>nigrum. Black pepper.</i>	5s.
—	<i>rugosum</i>	5s.
<i>Pistacia terebinthus. China Turpentine</i>	5s.
<i>Quisqualis indica</i>	10s.
<i>Ravenala madagascariensis. Urania</i>	5-25s.
<i>Rhus succedanea Japan Wax</i>	15s.
—	<i>vernicifera. Lacquer Tree</i>	10s.
<i>Saccharum officinarum. Sugar cane.</i>	2s.
—	<i>violaceum.</i>	5s.
—	<i>vittatum</i>	5s.
<i>Santalum album</i>	15s.
<i>Sapindus emarginatus</i>	15s.
—	<i>indica</i>	5s
—	<i>Saponaria. Soap Berry</i>	5s.
<i>Simaba Cedro</i>	30s.
<i>Simaruba amara</i>	30s.
—	<i>officinale</i>	15s.
<i>Smilax medica Sarsparilla</i>	15s.
—	<i>Salsaparilla</i>	10s.
<i>Stadmannia australis. Iron-wood</i>	5s.
—	<i>sideroxylon</i>	10s.
<i>Sterculia Chicha</i>	25s.
—	<i>Cola</i>	30s.

Stillingia sebifera. <i>Chinese Tallow Tree</i>	15s.
Strychnos Cabalonga	25s.
— nux vomica. <i>Nux Vomica</i>	25s.
Syncarpaea albens. <i>Turpentine Tree</i>	10s.
Tamarindus indica. <i>Tamarind</i>	5s.
Tecoma pentaphylla. <i>Roble blanco</i>	15s.
Terminalia Catappa. <i>Indian Almond</i>	10s.
Thea assamica. <i>Tea</i>	5s.
Theobroma Cacao. <i>Cocoa</i>	15s.
Thespesia populnea	10-20s.
Vangueria edulis. <i>Voa Vanga</i>	15s.
Vanilla aromatica. <i>Vanilla</i>	10s.
Wormia ferruginea	10s.
Xanthoxylon tragodes	15s.
Zingiber officinalis. <i>Ginger</i>	5s.

TROPICAL FRUIT TREES.

Achras dissecta	10s.
Anona Cherimolia. <i>Cherimoyer</i>	10s.
— — var. Loxensis	15s.
— muricata. <i>Soursop</i>	10s.
— reticulata. <i>Custard Apple</i>	10s.
— squamosa. <i>Sweet Sop</i>	10s.
Artocarpus grandis	30s.
— incisa. <i>Bread-fruit</i>	25s.
— integrifolia. <i>Jack-fruit</i>	10s.
— Neo Caledonica	15s.
Averrhoa Carambola. <i>Carambola</i>	5s.
Barringtonia racemosa (<i>Stravadium</i>)	10s.
— speciosa	15s.
Bertholetia excelsa. <i>Brazilian Nut</i>	25s.
Campomanesia hypoleuca.	15s.
Carica Candamarcensis	10s.
— erythrocarpa (ILLSTR.)	10s.
— Papaya. <i>Papaw tree</i>	3s.
Casimiroa edulis. <i>Mexican apple</i>	25s.
Chrysobalanus Iaco, var. Hondensis. <i>Cocoa Plum</i>	15s.
Chrysophyllum Cainito. <i>Star-apple</i>	5s.
Cicca disticha.	3s.
Cordia Sebestena	10s.
Crataeva gynandra. <i>Garlic Pear</i>	5s.
Diospyros embryopteris	5s.

Durio Zibethinus. <i>Durian</i>	25s.
Eriobotrya japonica. <i>Loquat</i>	1s.
Eugenia brasiliensis. (<i>Grumichama</i>)	5s.
— <i>Mitchelli (Pitanga)</i> .	5s.
Ficus macrocarpa	10s.
Fusanus acuminatus	10s.
Garcinia australis <i>Mangoosteen</i>	10s.
— <i>Livingstoni</i>	25s.
Jambosa magnifica	15s.
— <i>Malaccensis. Malay-Apple</i>	5s.
Lecythis Pisonis	15s.
Lucuma mammosa. <i>Marmalade-tree</i>	15s.
Malpighia punicifolia	10s.
Melicocca bijuga	10s.
— <i>olivæformis</i>	10s.
Mimusops cyanocarpa	5s.
— <i>Elengi. Medlar of Surinam.</i>	15s.
Musa Cavendishi. <i>Dwarf Banana</i>	5-15s.
— <i>coccinea</i>	15s.
— <i>Ensete</i>	2-5s.
— <i>glauca</i>	10s.
— <i>paradisiaca. Plantain</i>	10s.
— <i>superba</i>	5s.
— <i>textilis. Manilla Hemp</i>	25s.
— <i>vittata</i>	50-75s.
— <i>zebrina</i>	15s.
Omphalia triandra. <i>Hognut</i>	15s.
Passiflora edulis. <i>Granadilla</i>	5s.
— <i>macrocarpa</i>	5s.
— <i>quadrangularis. Granadilla</i>	2s.
Persea gratissima. <i>Alligator Pear</i>	10s.
Phoenix Dactylifera. <i>Date</i>	3s.
Physocalyx edulis.	10s.
Platonia granatensis	30s.
— <i>Insignis. Baçura-uva</i>	25s.
Sapota Achras. <i>Sapodilla Plum</i>	10s.
— <i>antioquensis</i>	15s.
Spondias Monbin	10s.
— <i>pleiogynne</i>	10s.
Tamarindus indica. <i>Tamarind</i>	5s.
Trichilia Spondioïdes	5s.
Vangueria edulis. <i>Voa Vanga</i>	10s.
Xanthochymus ovalifolius	15s.

CALADIUMS.

25, our selection	25s.
50, in 25 varieties our selection	50s.
100, in 50 " " "	150s.

Araucaria.

<i>Araucaria excelsa</i>	<i>from</i> 10-15-50s.
— — <i>aurea marginata</i>	" 15-25s.
— — <i>pendula</i>	" 10-25.
— — <i>viridis</i>	" 10-20.
— — <i>glauca</i>	" 10-15-50s.
— — <i>albo spica</i>	" 10-250s.
— — <i>Bidwillii</i>	" 10-50s.
— — <i>Cooki</i>	" 10s.
— — <i>Cunninghami</i>	" 10s.
— — <i>elegans</i>	" 15s.
— — <i>Rulei</i>	" 25s.
— — <i>multiceps</i>	" 15-25s.
— — <i>Mülleri</i>	" 40s.



CHOICE CAMELLIAS

FIGURED IN THE ILLUSTRATION HORTICOLE

Each : 2 shillings and 5 shillings according to size.

AT OUR OWN SELECTION :

12 different Camellias, figured in the ILLUSTRATION HORTICOLE, 20s.
The complete collection of 62 varieties ; 115s. to 200s., according to size.

- 1294. **Adriana.** Rose cerise; petals unusually large.
- 1162. **Alba ornatissima.** Pure white; perfectly imbricated.
- 1516. **Albino Botti.** An admirable plant with stout wood and handsome foliage; leaves acutely acuminate; distantly toothed. Flowers among the largest of the genus, regularly imbricated in zones with obcordate, flat, entire or slightly emarginate petals, beautiful deep rose in centre, encircled by a marginal band of pure white.

-
- 1354. **Angelo Cocchi.** White ground tinged with pink with broad crimson-purple stripes.
 - 1359 **Archiduc Étienne.** Transparent pink, spotted with bright crimson.
 - 1325. **Baron de Vrière.** A finely imbricated lovely pink flower, paler towards the centre.
 - 1229. **Bella Romana.** Resembling a huge flake carnation, pink striped with bright crimson.
 - 1437. **Bertha Giglioli.** Flower medium, delicate pink veined with rose; petals margined with white and perfectly imbricated.
 - 1316. **Caprioli.** Flower medium, regularly imbricated; petals oblong, deeply emarginate, of a beautiful bright crimson, each petal traversed longitudinally by a broad white stripe; the regularity of which gives the flower a starlike appearance.
 - 1383. **Carlotta Peloso.** Flower very large and perfectly imbricated bright rose, scarlet streaked with white.
 - 1430. **Caterina Rossi.** Delicate rose streaked with purple; perfectly imbricated.
 - 1343. **Clodia.** Flower large bright rosy red with distant white stripes.
 - 1326. **Comte de Toll.** Habit elegant; pink veined with carmine and striped with white.
 - 1303. **Comtesse Pasolini.** Clear delicate pink passing to white towards the margin. A free bloomer.
 - 1397. **Constantin Tretiakoff.** White ground, tinged with pink, more intense towards the centre.
 - 1438. **Contessa Tozzoni.** Flower regularly imbricated; beautiful bright rose; passing into almost pure white on the margin.
 - 1267. **Cora L. Barton.** Pure white ground, finely pencilled with pink.
 - 1293. **Dionisia Poniatowski.** Flower very large, white faintly shaded with delicate pink, with one or two streaks of crimson.
 - 1471. **Don Carlos Ferdinando.** Flowers very large and perfectly imbricated; petals obtuse scarcely emarginate arranged in radiating zones of a beautiful blood-red inclining to crimson, here and there with a slight dash of pure white near the tip.
 - 1473. **Don Pedro.** A first class variety with handsome regularly imbricated flowers; petals oval rounded, mucronate, distinct of good substance, beautiful pure white, longitudinally and faintly striped with pink here and there, orbicular-reniform in outline.
 - 1280. **Duchesse de Nassau.** Petals slightly cut and erose at the top; delicate pink on a white ground.
 - 1427. **Elvina Delli.** Flowers perfectly imbricated, delicate pink, of moderate size; petals margined with white and traversed with rose-coloured veins, a central longitudinally rosy band completing the ornamental effect of this admirable variety.
 - 1299. **Fanny Sanchioli.** Pure white, with a slight flush of yellow in the centre and scattered pink spots.
 - 1428. **Francesco Burlamachi.** Very regularly imbricated: petals broadly oval, slightly emarginate; the central ones irregular, oblong of a beautiful bright carmine finely striped, with a broad central white band expanding towards the summit and tinged with yellow in the centre.

1477. **Giardino Santarelli.** Flower of the largest size, and perfectly imbricated, bright cerise red, some of the petals half red and white, or wholly white with a crimson stripe.
1260. **Giardino Schmitz.** Delicate pink; petals broad and nicely imbricated.
1321. **Giuseppe Biasi.** Flower large, pure white irregularly streaked with delicate pink, well set.
1457. **Il Giojello.** Flower medium, well formed, regularly imbricated, with orbicular depressed emarginate obtuse petals of the most beautiful carmine red or crimson very finely striped with lines of a deeper hue.
1385. **Isabella Orsini.** Delicate pink, large flower.
1424. **Italia unita.** Large imbricated flower of a splendid scarlet, reticulated with a deeper tinge.
1462. **Japonica fol. var.**
1125. **Lavinia Maggi.** White tinged with pink elegantly striped and rayed with deep rose; perfect habit.
1458. **Luisa Bartolini.**
- 1468 **Madame Ambroise Verschaffelt.** Imbrication perfect, of a whitish slightly tinged very clear colour, veined and sometimes dotted with a rosy red tint. Habit exceedingly elegant.
1479. **Madame Cachet.** Good habit; flowers very large and reflexed; petals regularly imbricated orbicular, scarcely emarginate. Ground white with a delicate transparent shade of pink of great clearness. Some of the petals spotted near the margin or finely striped with rose lake of a more or less deep tint.
1476. **Madame de Cannart d'Hamale.** Large erect flowers, well expanded and regularly imbricated. Petals obtuse, emarginate, incurved at the apex, of the most delicate pink slenderly veined with a deeper tinge, and provided with a central white band, shaded with salmon, which divides the petals into two halves, giving the plant a very distinct and novel appearance.
1412. **Madame Rudolph Abel.** Flowers large rose-pink tinged with violet and undecided white streaks in the centre; imbrication perfect.
1413. **Marchesa Davia.** Large handsome perfectly formed flowers, with imbricated, reflexed orbicular or reniform petals of a bright crimson red with longitudinal regular very slender stripes; some of the petals striped or dotted with white in the centre.
1390. **Marianna Talenti.** Bright cerise red, veined with a deeper tint, and banded with a pure white.
1410. **Mateo Molfino.** Bright cerise, traversed by broad streaks of pure white.
1478. **Mistress Dombrain.** Delicate pink white towards the margin of the petals, and veined with a deeper tint.
1465. **Miss Minnie Merritt.**
1337. **Nazzari.** Flower imbricated cupshaded; red ground with broad bands of white; free bloomer.
1310. **Ninfa del Tebro.** Bright cerise, with broad bands of white forming six regular rays.
1323. **Petazzi.** Imbrication perfect; rose-cerise and white curiously intermingled.
1430. **Pietro Boutourlin.** Flower large perfectly imbricated without being too

- close; petals somewhat cordate spreading and slightly deflexed, of a violet red striped with a deeper shade, especially at the base; central petals striped and doted with white especially towards the top.
1319. **Planipetana.** White with a tinge of pale yellow; free flowering; imbrication perfect.
1399. **Poldina Vanturi.** A very large regularly imbricated flower. Petals orbicular, spreading, emarginate, of a bright carmine with deeper coloured striations.
1381. **Principessa Clotilde.** Flower of medium size; white and pink ground streaked with rose.
1355. **Reine des Beautés.** Delicate rose, brighter towards the centre.
1283. **Reticulata flore pleno.** A very large flower of a bright cerise red shaded with pink.
1418. **Roma Risorta.** Very large, bright rose speckled and dashed with crimson.
1286. **Sangalli.** White slightly tinted with delicate pink; imbrication perfect.
1463. **Sasanqua fol. var.**
1285. **Souvenir d'Emile Defresne.** Dazzling vermillion red beautifully streaked with white.
1166. **Stella polare.** Rose carmine, striped with white and pink, of a square oblong outline, forming a real star.
1426. **Terisita Canzio Garibaldi.**
1330. **Tricolor imbricata.** Beautifully variegated crimson on a pure white ground.
1336. **Vesillo dell' Arno.** White striped with delicate rose.
1327. **Vicomte de Nieuland.** Delicate rose, with a few faint white stripes near the centre.
1168. **Virginia Franco.** Pink and white ground with distant stripes or spots of a deeper tint.
1361. **Vittorio-Emmanuele II.** White ground tinged with pink, veined with carmine.

NEW CAMELLIAS

CAMELLIA MADAME LINDEN

(Illustration horticole)

This variety is a sport from *Madame Verschaffelt*, but it is much superior. The colour is more brilliant, the petals being of a delicate bright rose, each one being ornamented with a pale border which gives an effect of the greatest beauty to the flower.

Price 25s.

Camellia BARONNE DE BLEICHRÖDER

This is a novelty of the highest merit, and will be one of those most demanded.

The flowers are large, imbricated, snow-white, streaked with fine fire-red lines

Young plant.	10s.
Stronger.	15s. to 25s.



POPULAR

AZALEA INDICA

Only the best varieties are included in this selection.

A. Borsig. A very fine double pure white variety, of good form and great substance.

Alba illustrata fl. pl. Very fine double white.

Alice. Deepest rose, with large vermillion blotch; a magnificent double variety.

Amœna. Bright rosy purple, flowers small, hose-in-hose fashion, abundant bloomer, a very distinct and pleasing species, very hardy. Native of China.

Apollon. Pure white, sometimes lined with bright red.

Baron Edouard Osy. Bright orange, with amaranthe blotch, semi-double.

Belle Gantoise. Rosy salmon and white, very free.

Bernhard Andre. Dark violet-purple, semi-double, large and very showy.

— — alba. Very fine double white.

Bijou de Paris. White, striped with rose and rosy-salmon.

Cérès. White blotched with rose, a very fine bloomer..

Charles Leirens. Flowers large and good form, semi-double, very dark salmon.

Coloris nova. Carmine marked with rich brown.

Comtesse de Beaufort. Flowers of a very beautiful rich rosy colour, with a deep crimson blotch on the upper petals.

Comtesse de Flandre.

Comtesse Eugenie de Kerchove. Semi-double, white, flaked with reddish carmine.

Criterion. Salmon-pink, margined with white, spotted with red.

Daphne. Large white semi-double flowers, blotched with green and striped with red.

Dominique Vervaeene. Clear bright orange, double, fine shape.

Duc de Nassau. Rosy-purple, extra fine.

Duchesse Adelaide de Nassau. Splendid large flamed scarlet, fine variety.

Eclatante. Deep crimson, shaded rose, upper petals spotted.

- Empereur de Brésil.** Fine large double rose of exquisite form, banded with white, beautifully marked on the upper petals with glossy red.
- Etoile de Gand.** Flowers peach, deeply margined with white, spotted with crimson.
- Eugene Mazel.** Rosy-salmon, upper lobes violet.
- Ferdinand Kegeljan.** Flowers of good form and substance, light orange-salmon.
- Flag of Truce.** Large double white, fine form and substance, extra fine.
- Flambeau.** Glowing crimson, the deepest coloured Azalea yet raised, very bright and effective.
- François Devos.** Clear deep crimson, fine double.
- Gloire de Belgique.** White, striped with purple.
- Imbricata.** A most superb double white variety, occasionally flaked with rose.
- Iveryana.** Pure white, striped with rose; flowers large, fine form.
- Jean Vervaeene.** Rosy-salmon, spotted with deep crimson, edged with white.
- John Gould Veitch.** Flowers lilac-rose, netted and bordered with white, and blotched with saffron, a very fine variety.
- Juliette.** Superb rose, fine form, good habit.
- La Victoire.** Centre of the flower reddish, lighter towards the edges, beautifully spotted on the upper petals with crimson maroon.
- La superbe.** Fine form, colour rich lake, bordered with pale orange, spotted black.
- Madame Alex. Hardy.** Rosy-orange, speckled rose-violet spots; semi-double.
- Madame Camille Van Langenhove.** Pure white, sometimes striped, fine flower, double.
- Madame Charles Van Eeckhaute.** A first-class novelty, of the form and size, pure white, with the edges slightly crisped.
- Madame Dominique Vervaeene.** Soft rose, stained with white.
- Madame Iris Lefebvre.** Very double, dark orange, shaded with bright violet and blotched with chocolate, very fine.
- Madame Van Houtte.** White, richly flaked with rose and carmine, fine form and substance, and good habit, a great acquisition.
- Madame Vandercruyssen.** Bright clear rose in colour, of good form, semi-double.
- Mademoiselle Leonie Van Houtte.** Very fine white, flaked with rose, and spotted with sulphur yellow.
- Mademoiselle Louise de Kerchove.** Large snow-white festoons encircle the bright flesh-coloured flowers, flaked with orange, blotched finely with maroon, and shaded with carmine, semi-double.
- Mademoiselle Marie Lefebvre.** Good form and substance, pure white.
- Mademoiselle Marie Van Houtte.** Semi-double, white, flaked with rosy salmon.
- Marquis of Lorne.** The flower of finest form and substance and of a brilliant scarlet, a most desirable variety.
- Modele.** Bright rose, extra fine.
- Ovata.** A small flowering species with a profusion of flowers, cinnamon red.
- President Ghellinck De Walle.** Very large double, of an intense rose with blotch on upper petals, rayed with crimson.

- President Van den Hecke.** White, finely striped and speckled with bright crimson, centre yellow, a very pretty variety.
- Princess Alice.** Pure white, fine form and substance, extra fine.
- Princess Charlotte.** Large flower, rose, fine form.
- Princess Louise.** Flowers double, pure white, large and of good substance.
- Princess Stephanie Clothilde.** Double white fimbriated, sometimes striped.
- Punctata rosea.** Deep rose, fine for forcing.
- Punctulata.** Creamy white, striped with vermillion and scarlet.
- Raphael.** Pure white, double.
- Reine des Doubles.** Flowers clear rose, very double, fine.
- Reine des Pays-Bas.** Violet pink, margined with white, striped.
- Reine des Roses.** Beautiful bright rose, deeply spotted.
- Reine de Portugal.** Good form and substance, flowers double and pure violet.
- Roi de Hollande.** Colour dark, sanguineous red, spotted with black.
- Roi des Blancs.** Fine form and good substance.
- Roi Leopold.** Rich glossy crimson, fine form.
- Sigismund Rucker.** Flowers lilac rose, strongly netted and bordered with white, splendidly blotched with bright crimson.
- Sinensis.** Dark orange yellow, slightly spotted on the upper segments.
- Souvenir de Leon Maenhaut.** Bright rosy carmine.
- Souvenir de Prince Albert.** Warm peach rose, deeply margined with pure white, double flower, prolific bloomer, a most beautiful variety.
- Stella.** Bright orange scarlet, with rich violet upper segments.
- Vittata crispiflora.** White, shaded with purple and crimson, very free and fine for forcing.
- Vuurwerk.** Brilliant orange red.
- Wilson Saunders.** A very fine new single white variety, striped and blotched with vivid red.

AZALEA INDICA, 100 fine well-budded plants : 125s., 150s., 200s., 300s., 500s., 1000s., according to force.

*The largest selection of exhibition Azalea and Camellias
in the trade. Prices on application.*

HARDY GHENT AZALEAS

SELECT VARIETIES

12 fine sorts, our selection	18s. to 25s.
25 " " .	30s. to 40s.
50 " " .	60s. to 75s.
100 in 12 varieties. .	125s. 150s. to 200s.

AZALEA MOLLIS

SELECT NAMED VARIETIES

6 sorts well grown	15s.
12 " " " "	30s.
6 sorts stronger	25s. to 30s.
10 " " " "	50s.
100 in 12 varieties very strong	200s. 250s. 300s. 400s.

AZALEA MOLLIS

VARIETIES FROM SEED

Per hundred	180s. 200s. 250s.
-----------------------	-------------------

AZALEA MOLLIS

These brilliant novelties have gained the highest rewards at various shows and were figured in the *l'Illustration Horticole* in 1878.

Comte de Kerchove.
Albicans.
Comtesse de Kerchove.

Mirabilis.
Aurea floribunda.
Ambroise Verschaffelt.

The variety	4s.
The six	20s.

Carnation Mad^{lle} Else de Bleichröder

Red Variety of the " SOUVENIR DE LA MALMAISON "

ILLUSTRATION HORTICOLE, 1881

This first class novelty resembles its predecessor exactly in form and shape, the foliage being slightly more delicate.

The flower is of a deep pink hue and altogether forms one of the most distinct and striking varieties we have ever seen.

Each : 5s. — The half dozen : 20s. — The dozen : 35s.

Opinion of «The Gardeners' Chronicle»

THE PRINCIPAL ENGLISH HORTICULTURAL JOURNAL

on the GENERAL HORTICULTURAL SHOW opened by the

COMPAGNIE CONTINENTALE D'HORTICULTURE

at Ghent from the 16th to the 22nd April 1883.

Some idea of the magnitude of the business here carried on may be gained when we say that the space covered by glass is in round numbers 100,000 feet, exclusive of pits; that the heating arrangements include the use of 26,000 feet of hot-water pipes; and that it has lately acquired 50 acres in the Isle of Levant, in the Mediterranean, for the cultivation of Palms and Camellias in the open air, and where 25,000 Kentias are already planted. A handsome new entrance to the nursery is being made on the bank of the Coupure Canal, three minutes' walk from the Casino, and which will be formally opened on the 15th, when the whole establishment will be opened to visitors as an exhibition in itself. Entering the nursery from this point the first object of interest will be a fine new octagonal pavilion containing flowering Orchids. To the left of this the visitor of a mechanical turn will find a new engine driven by the modern force, electricity, and which is used solely for pumping the water required in the nursery. Near by is a very fine new span-roofed house, over 100 feet long, 40 feet wide, and 26 feet high in the centre, which will contain 500 specimen Indian Azaleas in full bloom. Palms, large and small, form the leading speciality of the nursery, and sixteen houses are devoted to their cultivation, five of which are filled with seedlings. Specimen Palms occupy much room, and magnificent in size and richness of foliage are many of the examples to be seen, notably in the large Palm-house or winter garden, which contains besides several notable specimens of *Dicksonia antarctica*, with stems straight as gun barrels, 25 feet high, and 7 feet in circumference, and almost as noteworthy are stems of *Cyathea medullaris robusta*, whose long black scaly hairs give the boles the appearance of being clothed in bear-skins. In other Palm-houses are handsome examples of the true *Thrinax barbadense*, a scarce species, *Pritchardia macrocarpa*, *Kentia Lindeni*, a species from New Caledonia, with wine-coloured stems and leaves; *Pritchardia pacifica*, a noble plant; and *P. grandis*, a handsome specimen, of medium size, and now believed to be the largest in Europe; *Areca nobilis*, a handsome spiny-stemmed species; *Phoenix Andersoni*, new and as yet scarce, in the way of *Cocos Weddelliana*, and very elegant. Ferns are abundant in all sizes, from the lords of the forest above-mentioned to the humble sporelings in thumb pots. Orchids occupy seven houses, and are well grown, especially the cool section, whose homemade bulbs are giants by comparison with the imported ones. Specimen Orchids are not grown to the same

extent as by some of our nurserymen, medium saleable sizes being the rule here, but we should not omit mention of a grand specimen of *Sobralia macrantha alba*, 3 feet through. One house is specially devoted to *Vandas*, another to *Pitcher-plants*, another to *Pandanads*, and yet another to *Dracænas*, *Alocasias*, and *Dieffenbachias* &c. Several houses contain "mixtures," and interspersed about these structures will be found many sterling novelties, which we must allude to on another occasion. A sight never before seen anywhere, and not soon to be forgotten, is a long, low span-roofed house, containing 300,000 seedling *Tillandsias* and *Massangeas*, of some twenty-two sorts, and all about 1 inch high. Indian *Azaleas* are present in thousands—many thousands, and as a new departure with this plant the company is entering into the cultivation of specimen sizes, and have a grand lot already which will be full bloom during the show week. We have not space on this occasion to note a tithe of the interesting features of this fine establishment, and must defer further remarks till another occasion.

Seldom in our recollection has a nursery establishment been seen in such perfect show order as was that of the Compagnie Continentale d'Horticulture at Ghent, which was visited by H. M. the Queen of the Belgians on April 15, and opened for the inspection of visitors during the following week. An imposing new entrance from the Coupure has recently been constructed, thus allowing of a direct communication between that thoroughfare and the Rue du Chaume. As—for reasons which it is to be hoped will not be infectious—the company did not show in the general exhibition, or only to a limited extent, it is incumbent on us to advert thus separately to what was indeed a very beautiful and a very remarkable display. Everything about the place was spick and span, and the display of Indian *Azaleas* and other flowering plants simply marvellous. The new show-house, alluded to in a previous number, was a floral picture not soon to be forgotten, being filled entirely with large and small specimen *Azaleas*, grouped with the most perfect good taste in irregular shaped beds, and relieved to some extent by "dot" plants of handsome specimen Palms and Tree Ferns. Near by was a long, narrow spanrooofed house filled with half-specimen standard *Azaleas*, all perfectly flowered, and as if cast in the same mould. The plants were arranged in four rows, two on either side of the path, with light and dark coloured varieties alternating, and the result was a vista of colour surprisingly beautiful to the eye of a "practical." The large winter garden had been re-arranged and thinned out so as to allow of the unique specimens of Tree Ferns and Palms being seen to greater advantage, and many were the admirers of the highly tropical character of the *tout ensemble*. The new octagonshaped pavilion contained a choice assortment of Orchids, and other select Orchids in flower were to be seen in the other houses. The show of *Camellias*, though so late, was very good. Each of the great number of houses contained some special type or section of plants, all full to overflowing, and all arranged with a view to producing pictorial effect.

Among the new or less known plants, to some of which we have already alluded as having been exhibited, we may mention *Aralia ? Deleauana*, Hort., a form from the Solomon Isles, with multiform much divided glabrous leaves of a dark green colour. In some cases the ultimate segments are almost linear, in others obovate or roundish and coarsely toothed, so that the whole leaf reminds one, in

some instances, of the Parsley Fern (*Allosorus*), in others of an *Adiatum* like *A. Ludde-mannianum*. *Begonia Meysseliana*, Hort., is a Sumatran form, with obliquely oblong dull olivegreen leaves spotted with white. It is recommended for summer bedding. *Calamus? Lindenii*, Rodigas, has the stem covered with brownish down, from amid which project numerous short spines. The general outline of the pinnate leaves is oblong-obtuse, the pinnæ oblong acute, dark green above, white beneath, and minutely spiny at the edge. It is a very distinct and handsome Palm. *Vriesia bellula*, Hort., is a dwarf Bromeliad, with a tuft of purplish leaves, from the centre of which uprises a short spike of inflorescence consisting of two rows of boat-shaped bracts, red at the base, yellowish-green above, and which subtend small white flowers. *Aralia reginæ*, Hort. Linden — for our illustration of which (fig. 88, p. 564) we are indebted to M. Lucien Linden — is a New Caledonian form, the stem and leafstalks of which have that peculiar pinkish-olive hue, dotted with darker spots, common to so many Araliads. The blades of the leaves are very deeply divided, in palmate fashion, into 5—9 linear acute deep green segments. The plant was figured in the *Illustration horticole* for February, 1879. *Heliconia triumphans*, Hort. Lind. (fig. 89, p. 565), is a Sumatran form, with oblong acute leaves, dark green picked out with narrow blackish stripes running from the midrib to the margin at distant intervals. *Dracæna Lindenii* is a very striking plant, with narrow oblong recurved deep green leaves with one or more broad central stripes of golden-yellow. *Aglaonema pictum* is an Aroid with shortly stalked sheathing leaves, with ovate lanceolate blades of a rich green colour, marbled and mottled with white, especially along the midrib. *Begonia Diadema*, Hort. Lind., has obliquely ovate leaves, deeply and irregularly lobed, and of a shining green colour, mottled with milky-white spots. *Dieffenbachia magnifica* is a tall-growing Aroid, introduced from Venezuela, with acuminate leaves of a clear green colour, thickly sprinkled with whitish blotches, giving the appearance as if a slight shower of snow had fallen on the leaves.

Of a new and very distinct *Cattleya*, *C. nobilior*, we hope to give an illustration on another occasion. Perhaps the most astonishing sight in the nursery — at any rate most unusual to our eyes — was the prodigious quantity of seedling Bromeliads, Tillandsias, etc., in the propagating-houses, hundreds of square pans filled with seedlings of various sizes and stages of growth, but all arranged with such symmetry and regularity that if they had been pins in a cushion, disposed at regularly marked intervals, they could not have been more evenly arranged. What can be the destiny of all these millions seedlings? There are, as might be expected, considerable variations, according to the species; and M. Linden would be doing a service to science if he would publish in the *Illustration* a description of the seedling plant of each species.

(*Gardeners' Chronicle*).

